arXiv:0907.2935v1 [math.DS] 16 Jul 2009

Positive expansiveness versus network dimension in symbolic dynamical systems

Marcus Pivato
Department of Mathematics, Trent University, Canada

July 16, 2009

Abstract

A 'symbolic dynamical system' is a continuous transformation $\Phi : \mathcal{X} \longrightarrow \mathcal{X}$ of closed perfect subset $\mathcal{X} \subseteq \mathcal{A}^{\mathbb{V}}$, where $\mathcal{A}$ is a finite set and $\mathbb{V}$ is countable. (Examples include subshifts, odometers, cellular automata, and automaton networks.) The function Φ induces a directed graph structure on $\mathbb{V}$, whose geometry reveals information about the dynamical system $(\mathcal{X}, \Phi)$. The 'dimension' $\dim(\mathbb{V})$ is an exponent describing the growth rate of balls in the digraph as a function of their radius. We show: if $\mathcal{X}$ has positive entropy and $\dim(\mathbb{V}) > 1$, and the system $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ satisfies minimal symmetry and mixing conditions, then $(\mathcal{X}, \Phi)$ cannot be positively expansive; this generalizes a well-known result of Shereshevsky about multidimensional cellular automata. We also construct a counterexample to a version of this result without the symmetry condition. Finally, we show that network dimension is invariant under topological conjugacies which are Hölder-continuous.

Let $\mathcal{X}$ be *Cantor space* (the compact, perfect, zero-dimensional metrizable topological space, which is unique up to homeomorphism). A *Cantor dynamical system* is a continuous self-map $\Phi : \mathcal{X} \longrightarrow \mathcal{X}$. In addition to its intrinsic interest, the class of Cantor systems is important because it has two universal properties. First, any topological dynamical system on a compact metric space is a factor of a Cantor system; see [Kůr03, Corollary 3.9, p.106] or [BS89, p.1241]. Second, the Jewet-Krieger Theorem says that any ergodic measure-preserving system can be represented as a uniquely ergodic, minimal Cantor system [Pet89, §4.4, p.188].

If $\mathcal{A}$ is a finite set, and $\mathbb{V}$ is a countably infinite set, then the product space $\mathcal{A}^{\mathbb{V}}$ is a Cantor space. Thus, any Cantor dynamical system can be represented as a self-map $\Phi : \mathcal{A}^{\mathbb{V}} \longrightarrow \mathcal{A}^{\mathbb{V}}$, or more generally, as a self-map $\Phi : \mathcal{X} \longrightarrow \mathcal{X}$, where $\mathcal{X} \subset \mathcal{A}^{\mathbb{V}}$ is a *pattern space* (a closed perfect subset of $\mathcal{A}^{\mathbb{V}}$). We refer to the structure $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ as a *symbolic dynamical system*. At an abstract topological level, any pattern space $\mathcal{X}$ is homeomorphic to Cantor space, so a symbolic dynamical system is simply a Cantor dynamical system. What distinguishes symbolic dynamical systems is a particular way of representing $\mathcal{X}$ as a subset of some Cartesian product $\mathcal{A}^{\mathbb{V}}$ (so that an element of $\mathcal{X}$ corresponds to some $\mathbb{V}$-indexed 'pattern' of 'symbols' in the alphabet $\mathcal{A}$).

The *network* of Φ is the digraph structure ($\bullet\!\!\!\to$) on $\mathbb{V}$ defined as follows: for all $\mathsf{v}, \mathsf{w} \in \mathbb{V}$, we have $\mathsf{v} \bullet\!\!\!\to \mathsf{w}$ if and only if the value of $\Phi(\mathbf{x})_{\mathsf{w}}$ depends nontrivially on the value of x_{v}. We say that $(\mathbb{V}, \bullet\!\!\!\to)$ has *dimension* δ if the cardinality of a ball of radius r grows like r^{δ} as $r \rightarrow \infty$. (Note that δ is not necessarily an integer.) For example, if $\Phi : \mathcal{A}^{\mathbb{Z}^D} \longrightarrow \mathcal{A}^{\mathbb{Z}^D}$ is a cellular automaton, then its network is just a Cayley digraph on $\mathbb{Z}^D$; the dimension of this network is D.

This paper explores the relationship between network dimension and the properties of $(\mathcal{X}, \Phi)$ as a topological dynamical system. In §1, we formally define the dimension of a network $(\mathbb{V}, \bullet\!\!\!\to)$ and the entropy of a pattern space on $\mathbb{V}$. In §2, we generalize an important result of Shereshevsky (later reproved by Finelli, Manzini, and Margara) about multidimensional cellular automata. We show: if $\dim(\mathbb{V}, \bullet\!\!\!\to) > 1$ (more generally, if $(\mathbb{V}, \bullet\!\!\!\to)$ has 'superlinear connectivity'), and $\mathcal{X}$ has positive entropy and a mild 'mixing' condition, and the system $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ has some minimal symmetries, then $(\mathcal{X}, \Phi)$ cannot be positively expansive (Theorem 2.7). In §3, we consider the *propagation* of a symbolic dynamical system, and its relationship with sensitivity and equicontinuity. In §4, we show that a 'naive' generalization of Shereshevsky's result cannot be true, by constructing a positively expansive symbolic dynamical system with network dimension two. Thus, any result similar to Theorem 2.7 must impose at least some additional technical conditions.

The counterexample in §4 also shows that network dimension is not invariant under topological conjugacy; thus, it cannot be treated as a structural property of an abstract Cantor dynamical system $(\mathcal{X}, \Phi)$. However, in §5, we propose to augment the system $(\mathcal{X}, \Phi)$ with a metric which is Lipschitz for Φ; we show that network dimension *is* a structural property of this 'metric' Cantor system, as it is invariant under all biHölder conjugacies (Corollary 5.8). Sections 2-5 are logically independent, and can be read in any order.

1 Preliminaries

Let $\mathcal{A}$ be a finite set (called an *alphabet*) endowed with the discrete topology. Let $\mathbb{V}$ be a countably infinite set of points (called *vertices*). Endow $\mathcal{A}^{\mathbb{V}}$ with the Tychonoff product topology. A *pattern space* is a closed perfect subset $\mathcal{X} \subseteq \mathcal{A}^{\mathbb{V}}$. A *symbolic dynamical system* is triple $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$, where $\mathcal{X} \subseteq \mathcal{A}^{\mathbb{V}}$ is a pattern space and $\Phi : \mathcal{A}^{\mathbb{V}} \longrightarrow \mathcal{A}^{\mathbb{V}}$ is a continuous function such that $\Phi(\mathcal{X}) \subseteq \mathcal{X}$. (Sometimes we will simply indicate this as $(\mathcal{X}, \Phi)$ when $\mathcal{A}$ and $\mathbb{V}$ are clear from context.)

Example 1.1. (a) Let $\mathbb{V} = \mathbb{Z}^D \times \mathbb{N}^E$ for some $D, E \geq 0$ and let $\mathcal{X} \subset \mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E}$ be a *subshift* (i.e. closed, shift-invariant subset). Then $\mathcal{X}$ is a pattern space. Fix $\mathsf{z} \in \mathbb{Z}^D \times \mathbb{N}^E$, and let $\sigma^{\mathsf{z}} : \mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E} \longrightarrow \mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E}$ be the associated shift map. Then $(\mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E}, \mathcal{X}, \sigma^{\mathsf{z}})$ is a symbolic dynamical system.

(b) Let $\mathbb{V} = \mathbb{Z}^D \times \mathbb{N}^E$, and let $\Phi : \mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E} \longrightarrow \mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E}$ be a *cellular automaton* (CA) —i.e. a continuous, shift-commuting map. Then $(\mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E}, \Phi)$ is a symbolic dynamical system. More generally, if $\mathcal{X} \subset \mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E}$ is any Φ-invariant subshift, then $(\mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E}, \mathcal{X}, \Phi)$ is a symbolic dynamical system.

(c) For all $\mathsf{v} \in \mathbb{V}$, let $\mathcal{A}_{\mathsf{v}} \subseteq \mathcal{A}$. Let $\mathcal{X} := \prod_{\mathsf{v} \in \mathbb{V}} \mathcal{A}_{\mathsf{v}}$; then $\mathcal{X}$ is a pattern space. If $\Phi : \mathcal{X} \longrightarrow \mathcal{X}$ is a continuous self-map, then $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ is a symbolic dynamical system, sometimes called an ***automaton network***, because it can be interpreted as an infinite network of interacting finite-state automata.

(d) Gromov has initiated a study of 'proalgebraic' dynamical systems, which are (loosely speaking) projective limits of polynomial self-mappings of algebraic varieties [Gro99]. If the base field $\mathbb{F}$ is finite (e.g. $\mathbb{F} = \mathbb{Z}_{/p}$), then a 'proalgebraic space' can be represented as a pattern space with alphabet $\mathbb{F}$; hence a proalgebraic system is a symbolic dynamical system. ♢

The analysis of subshifts and cellular automata depends heavily on the highly symmetric structure created by shift-invariance. Likewise, Gromov's analysis of proalgebraic systems requires a structure of 'local' symmetries (called *holonomies*). We will also make use of some minimal symmetry assumptions in §2. However, in general, symbolic dynamical systems do not have any appreciable symmetries.

For any $\mathbf{x} \in \mathcal{X}$ and $\mathbb{U} \subset \mathbb{V}$, we define $\mathbf{x}_{\mathbb{U}} := [x_{\mathsf{u}}]_{\mathsf{u}\in\mathbb{U}} \in \mathcal{A}^{\mathbb{U}}$; we then define $\mathcal{X}_{\mathbb{U}} := \{\mathbf{x}_{\mathbb{U}} \ ; \ \mathbf{x} \in \mathcal{X}\} \subseteq \mathcal{A}^{\mathbb{U}}$. A function $\gamma : \mathcal{A}^{\mathbb{U}} \longrightarrow \mathcal{B}$ is ***proper*** if γ depends nontrivially on every coordinate in $\mathbb{U}$. Formally: for every $\mathsf{v} \in \mathbb{U}$, there exist $\mathbf{a}, \mathbf{a}' \in \mathcal{A}^{\mathbb{U}}$ such that: (1) $a_{\mathsf{u}} = a'_{\mathsf{u}}$ for all $\mathsf{u} \in \mathbb{U} \setminus \{\mathsf{v}\}$; (2) $a_{\mathsf{v}} \neq a'_{\mathsf{v}}$; and (3) $\gamma(\mathbf{a}) \neq \gamma(\mathbf{a}')$.

Lemma 1.2 *Let* $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ *be a symbolic dynamical system. Then for all* $\mathsf{v} \in \mathbb{V}$*, there is some finite subset* $\mathbb{U}(\mathsf{v}) \subset \mathbb{V}$ *and a proper function* $\phi_{\mathsf{v}} : \mathcal{A}^{\mathbb{U}(\mathsf{v})} \longrightarrow \mathcal{A}$ *such that, for any* $\mathbf{x} \in \mathcal{X}$*,* $\Phi(\mathbf{x})_{\mathsf{v}} = \phi_{\mathsf{v}}(\mathbf{x}_{\mathbb{U}(\mathsf{v})})$.

Proof. For each $\mathsf{v} \in \mathbb{V}$, the existence of a local rule ϕ_{v} is proved by exactly the same argument as the Curtis-Hedlund-Lyndon theorem for cellular automata (see e.g. [Kůr03, Theorem 5.2, p.190]; observe that the construction of the local rule does not depend on shift-invariance). The difference is that there may be different local rules at different vertices. □

The set $\mathbb{U}(\mathsf{v})$ is called the ***input neighbourhood*** of Φ at v, and denoted $\Phi_{\text{in}}(\mathsf{v})$. The function $\phi_{\mathsf{v}} : \mathcal{A}^{\Phi_{\text{in}}(\mathsf{v})} \longrightarrow \mathcal{A}$ is called the ***local rule*** of Φ at w.

Example 1.3. (a) If $\Phi : \mathcal{A}^{\mathbb{Z}^D} \longrightarrow \mathcal{A}^{\mathbb{Z}^D}$ is a cellular automaton, then Corollary 1.2 plus shift-invariance yields the Curtis-Hedlund-Lyndon theorem.

(b) Fix $A \in \mathbb{N}$ and let $\mathcal{A} := [0...A)$. Let $\mathbf{m} := (m_0, m_1, m_2, \ldots)$ be a sequence of natural numbers in $[1...A]$. Let $\mathbb{V} := \mathbb{N}$, and let $\mathcal{X} := \{\mathbf{a} \in \mathcal{A}^{\mathbb{N}}; \ 0 \leq a_{\mathsf{v}} < m_{\mathsf{v}}, \ \forall \ \mathsf{v} \in \mathbb{V}\}$. Let $\Phi : \mathcal{X} \longrightarrow \mathcal{X}$ be the $\mathbf{m}$-ary odometer [Kůr03, §4.1, p.136]. Then $\Phi_{\text{in}}(0) = \{0\}$ and $\phi_0 : \mathcal{A} \longrightarrow \mathcal{A}$ is defined by $\phi_0(a_0) := (a_0 + 1) \bmod m_0$. Meanwhile, for all $N \geq 1$, we have $\Phi_{\text{in}}(N) = [0...N]$, and $\phi_N : \mathcal{A}^{[0...N]} \longrightarrow \mathcal{A}$ is defined by

$$\phi_N(a_0, a_1, \ldots, a_N) \quad := \quad \begin{cases} (a_N + 1) \bmod m_N & \text{if } \ a_n = m_n - 1, \ \forall n \in [0...N); \\ \qquad a_N & \text{otherwise.} \end{cases} \qquad \diamondsuit$$



Figure 1: The network of an odometer.

Directed graphs. Let $\mathbb{V}$ be a set of 'vertices'. A *directed graph* (or *digraph*) structure on $\mathbb{V}$ is a binary relation $(\bullet\!\!\to)$ on $\mathbb{V}$ (i.e. a subset $(\bullet\!\!\to) \subseteq \mathbb{V} \times \mathbb{V}$). For any $\mathsf{v}, \mathsf{w} \in \mathbb{V}$, we write "$\mathsf{v} \bullet\!\!\to \mathsf{w}$" if $(\mathsf{v}, \mathsf{w}) \in (\bullet\!\!\to)$. More generally, we say v is *upstream* of w ("$\mathsf{v} \rightsquigarrow \mathsf{w}$") if either $\mathsf{v} = \mathsf{w}$, or there is a directed path $\mathsf{v} = \mathsf{v}_0 \bullet\!\!\to \mathsf{v}_1 \bullet\!\!\to \cdots \bullet\!\!\to \mathsf{v}_n = \mathsf{w}$. The relation $(\rightsquigarrow)$ is a partial order (it is reflexive and transitive). We write $\mathsf{v} \leftrightsquigarrow \mathsf{w}$ if $\mathsf{v} \rightsquigarrow \mathsf{w}$ and $\mathsf{w} \rightsquigarrow \mathsf{v}$. Thus, $(\leftrightsquigarrow)$ is an equivalence relation; the $(\leftrightsquigarrow)$-equivalence classes of $\mathbb{V}$ are called the *biconnected components* of $(\mathbb{V}, \bullet\!\!\to)$. We say that $(\mathbb{V}, \bullet\!\!\to)$ is *biconnected* if $\mathsf{v} \leftrightsquigarrow \mathsf{w}$ for all $\mathsf{v}, \mathsf{w} \in \mathbb{V}$.

Let $(\sim)$ be the smallest equivalence relation on $\mathbb{V}$ which contains $(\bullet\!\!\to)$. Equivalently, for any $\mathsf{u}, \mathsf{w} \in \mathbb{V}$, we have $\mathsf{u} \sim \mathsf{w}$ if either (1) $\mathsf{u} \rightsquigarrow \mathsf{w}$; or (2) $\mathsf{w} \rightsquigarrow \mathsf{u}$; or (3) (inductively) there exists some $\mathsf{v} \in \mathbb{V}$ such that $\mathsf{u} \sim \mathsf{v} \sim \mathsf{w}$. The $(\sim)$-equivalence classes are the *connected components* of $\mathbb{V}$; if $\mathsf{v} \sim \mathsf{w}$ for all $\mathsf{v}, \mathsf{w} \in \mathbb{V}$, then we say that $(\mathbb{V}, \bullet\!\!\to)$ is *connected*.

If $\Phi : \mathcal{A}^{\mathbb{V}} \longrightarrow \mathcal{A}^{\mathbb{V}}$ is any continuous function, then we can define a digraph relation $(\bullet\!\!\to)$ on $\mathbb{V}$ by $\Big(\mathsf{v} \bullet\!\!\to \mathsf{w}\Big) \Leftrightarrow \Big(\mathsf{v} \in \Phi_{\mathrm{in}}(\mathsf{w})\Big)$, where $\Phi_{\mathrm{in}}(\mathsf{w})$ is as defined by Lemma 1.2 above. This digraph is called the *network* of Φ.

Example 1.4. (a) Let $\Phi : \mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E} \longrightarrow \mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E}$ be a cellular automaton; then the network of Φ is a Cayley digraph on $\mathbb{Z}^D \times \mathbb{N}^E$.

(b) Figure 1 depicts the network of the odometer $\Phi : \mathcal{A}^{\mathbb{N}} \longrightarrow \mathcal{A}^{\mathbb{N}}$ from Example 1.3(b). $\diamondsuit$

Connectivity dimension. Let $(\mathbb{V}, \bullet\!\!\to)$ be an infinite digraph (e.g. the network of a continuous function $\Phi : \mathcal{A}^{\mathbb{V}} \longrightarrow \mathcal{A}^{\mathbb{V}}$). For any subset $\mathbb{U} \subset \mathbb{V}$, define $\mathbb{B}(\mathbb{U}, 1) := \mathbb{U} \cup \{\mathsf{v} \in \mathbb{V} \; ; \; \exists \ \mathsf{u} \in \mathbb{U} : \ \mathsf{v} \bullet\!\!\to \mathsf{u}\}$. Then inductively define $\mathbb{B}(\mathbb{U}, n+1) := \mathbb{B}[\mathbb{B}(\mathbb{U}, n), 1]$ for all $n \in \mathbb{N}$. Thus, $\mathbb{B}(\mathsf{w}, 1) := \{\mathsf{w}\} \cup \Phi_{\mathrm{in}}(\mathsf{w})$, and $\mathbb{B}(\mathsf{w}, r)$ is the set of all $\mathsf{v} \in \mathbb{V}$ such that there exists some path $\mathsf{v} = \mathsf{v}_1 \bullet\!\!\to \mathsf{v}_2 \bullet\!\!\to \cdots \bullet\!\!\to \mathsf{v}_s = \mathsf{w}$ with $s \leq r$. For any $\mathsf{v} \in \mathbb{V}$, we define

$$\underline{\dim}_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\to) \ := \ \liminf_{r \to \infty} \frac{\log |\mathbb{B}(\mathsf{v}, r)|}{\log(r)} \quad \text{and} \quad \overline{\dim}_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\to) \ := \ \limsup_{r \to \infty} \frac{\log |\mathbb{B}(\mathsf{v}, r)|}{\log(r)}. \quad (1)$$

If $\underline{\dim}_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\to) = \overline{\dim}_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\to)$, then we refer to their common value as "$\dim_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\to)$", the *connectivity dimension* of $(\mathbb{V}, \bullet\!\!\to)$ at v, and we say that $(\mathbb{V}, \bullet\!\!\to)$ is *dimensionally regular* at v.

Example 1.5. For all $r \in \mathbb{N}$, let $\beta_{\mathsf{v}}(r) := |\mathbb{B}(\mathsf{v}, r)|$.

(a) Let $\delta \in [0, \infty)$, and suppose

$$0 \quad < \quad \liminf_{r \to \infty} \frac{\beta_{\mathsf{v}}(r)}{r^\delta} \quad \leq \quad \limsup_{r \to \infty} \frac{\beta_{\mathsf{v}}(r)}{r^\delta} \quad < \quad \infty.$$

(For example, suppose $\beta_{\mathsf{v}}(r) = C\, r^\delta + p(r)$, where C is a constant and p is a polynomial of degree less than δ.) Then $\dim_{\mathsf{v}}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}) = \delta$.

(b) Likewise, if $C, \delta, \lambda > 0$, and $\beta_{\mathsf{v}}(r) = C\, r^\delta \cdot \log(r)^\lambda$, then $\dim_{\mathsf{v}}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}) = \delta$.

(c) Let $c > 0$. If $\beta_{\mathsf{v}}(r) = c^r$, then $\dim_{\mathsf{v}}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}) = \infty$. ♢

Let $\overline{\dim}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}) := \sup \left\{ \overline{\dim}_{\mathsf{v}}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})\ ;\ \mathsf{v} \in \mathbb{V} \right\}$ and $\underline{\dim}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}) := \inf \left\{ \underline{\dim}_{\mathsf{v}}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})\ ;\ \mathsf{v} \in \mathbb{V} \right\}$. If $\underline{\dim}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}) = \overline{\dim}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$, then we refer to their common value as "$\dim(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$", the (global) *connectivity dimension* of $(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$, and we say that $(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$ is *dimensionally homogeneous*. (This implies that $(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$ is everywhere dimensionally regular.)

Example 1.6. (a) If $\mathbb{Z}^D$ has the obvious Cayley digraph structure, then $\dim(\mathbb{Z}^D) = D$.

(b) Let $(\mathbb{N}, \mathrel{\bullet\!\!\rightarrow})$ be the digraph in Figure 1 (the odometer). Then $\dim(\mathbb{N}, \mathrel{\bullet\!\!\rightarrow}) = 0$, because for all $n \in \mathbb{N}$, and all $r \geq 1$, we have $|\mathbb{B}(n, r)| = n + 1$ (because $\mathbb{B}(n, r) = [0...n]$).

(c) If $(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$ is the Cayley digraph of a group $\mathcal{G}$, then $\dim(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$ is the 'growth dimension' of $\mathcal{G}$. More generally, if $(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$ is any graph whose automorphism group acts transitively, then $(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$ is 'almost' a Cayley digraph; for a survey of the well-developed dimension theory for such graphs, see [MW89, §5] or [IS91]. ♢

The dimension of a Cayley digraph is always an integer. However, there exist 'self-similar' graphs with fractional connectivity dimensions [McC91]. Connectivity dimension is closely related to properties of diffusion processes and electrical conductance on graphs [Tel89, Tel90, Tel95, Tel01], the existence of periodic points in 'majority vote' networks [Mor95, GH00], and also arises in certain models of quantum gravity [NR99, NR98].

Not all digraphs are dimensionally regular. For example, consider a digraph which consists of increasingly large 'clumps' which are spaced at increasingly long intervals along an infinite line-graph; by making the clumps and the intervals between them grow fast enough, one can force $\underline{\dim}_{\mathsf{v}}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}) = 1$ while $\overline{\dim}_{\mathsf{v}}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}) > 1$ for some $\mathsf{v} \in \mathbb{V}$. (However, examples like this are highly contrived; probably, most 'natural' examples are dimensionally regular.)

Furthermore, not all connected, dimensionally regular digraphs are dimensionally homogeneous. For example, let $\mathbb{V}_1 \cong \mathbb{Z}$ be a biconnected Cayley digraph of $\mathbb{Z}$, and let $\mathbb{V}_2 \cong \mathbb{Z}^2$ be a biconnected Cayley digraph of $\mathbb{Z}^2$. Let $\mathbb{V} = \mathbb{V}_1 \sqcup \mathbb{V}_2$, with connections $n \mathrel{\bullet\!\!\rightarrow} (n, 0)$ for all $n \in \mathbb{Z} \cong \mathbb{V}_1$. Then $\mathbb{V}_1$ and $\mathbb{V}_2$ are biconnected components of $\mathbb{V}$, with $\mathbb{V}_1$ upstream from $\mathbb{V}_2$. Clearly, $\dim_{\mathsf{v}}(\mathbb{V}_k) = k$ for all $\mathsf{v} \in \mathbb{V}_k$ and $k = 1, 2$.

Lemma 1.7 *Let $(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$ be a digraph. If $\mathsf{v} \rightsquigarrow \mathsf{w}$, then $\underline{\dim}_{\mathsf{v}}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}) \leq \underline{\dim}_{\mathsf{w}}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$ and $\overline{\dim}_{\mathsf{v}}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}) \leq \overline{\dim}_{\mathsf{w}}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$.*

Proof. If $\mathsf{v} \rightsquigarrow \mathsf{w}$, then there exists $R > 0$ such that $\mathsf{v} \in \mathbb{B}(\mathsf{w}, R)$. Thus, for all $r \in \mathbb{N}$, we have $\mathbb{B}(\mathsf{v}, r) \subseteq \mathbb{B}(\mathsf{w}, R + r)$, hence $|\mathbb{B}(\mathsf{v}, r)| \leq |\mathbb{B}(\mathsf{w}, R + r)|$. Thus

$$\begin{aligned}
\underline{\dim}_{\mathsf{v}}(\mathbb{V}, \mathrel{\bullet\!\!\to}) &:= \liminf_{r\to\infty} \frac{\log|\mathbb{B}(\mathsf{v}, r)|}{\log(r)} \quad\leq\quad \liminf_{r\to\infty} \frac{\log|\mathbb{B}(\mathsf{w}, R + r)|}{\log(R + r)} \cdot \frac{\log(R + r)}{\log(r)} \\
&= \left(\liminf_{r\to\infty} \frac{\log|\mathbb{B}(\mathsf{w}, R + r)|}{\log(R + r)}\right) \cdot \left(\lim_{r\to\infty} \frac{\log(R + r)}{\log(r)}\right) \quad=\quad \underline{\dim}_{\mathsf{w}}(\mathbb{V}, \mathrel{\bullet\!\!\to}) \cdot 1.
\end{aligned}$$

Hence $\underline{\dim}_{\mathsf{v}}(\mathbb{V}, \mathrel{\bullet\!\!\to}) \leq \underline{\dim}_{\mathsf{w}}(\mathbb{V}, \mathrel{\bullet\!\!\to})$. Likewise, $\overline{\dim}_{\mathsf{v}}(\mathbb{V}, \mathrel{\bullet\!\!\to}) \leq \overline{\dim}_{\mathsf{w}}(\mathbb{V}, \mathrel{\bullet\!\!\to})$. ☐

If $\mathbb{W} \subset \mathbb{V}$ is a biconnected component of $(\mathbb{V}, \mathrel{\bullet\!\!\to})$, then Lemma 1.7 says that every vertex in $\mathbb{W}$ has the same connectivity dimension. In particular, if $(\mathbb{V}, \mathrel{\bullet\!\!\to})$ is biconnected and dimensionally regular, then it is dimensionally homogeneous.

Entropy. Let $(\mathbb{V}, \mathrel{\bullet\!\!\to})$ be a digraph, and let $\mathcal{X} \subset \mathcal{A}^{\mathbb{V}}$ be a pattern space. For any $\mathsf{v} \in \mathbb{V}$, we define the *lower* and *upper topological entropies* of $\mathcal{X}$ around v by:

$$\underline{h}_{\mathsf{v}}(\mathcal{X}) \quad:=\quad \liminf_{r\to\infty} \frac{\log_2|\mathcal{X}_{\mathbb{B}(\mathsf{v},r)}|}{|\mathbb{B}(\mathsf{v}, r)|} \qquad\text{and}\qquad \overline{h}_{\mathsf{v}}(\mathcal{X}) \quad:=\quad \limsup_{r\to\infty} \frac{\log_2|\mathcal{X}_{\mathbb{B}(\mathsf{v},r)}|}{|\mathbb{B}(\mathsf{v}, r)|}. \tag{2}$$

Clearly, $0 \leq \underline{h}_{\mathsf{v}}(\mathcal{X}) \leq \overline{h}_{\mathsf{v}}(\mathcal{X}) \leq \log_2|\mathcal{A}|$. Let $\underline{h}(\mathcal{X}) := \inf_{\mathsf{v}\in\mathbb{V}} h_{\mathsf{v}}(\mathcal{X})$ and $\overline{h}(\mathcal{X}) := \sup_{\mathsf{v}\in\mathbb{V}} h_{\mathsf{v}}(\mathcal{X})$.

2 Positive expansion versus network connectivity

An abstract Cantor dynamical system $(\mathcal{X}, \Phi)$ is *posexpansive* if it is topologically conjugate to a one-sided shift. In particular, let $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ be a symbolic dynamical system. Fix a finite subset $\mathbb{W} \subset \mathbb{V}$, let $\mathcal{B} := \mathcal{A}^{\mathbb{W}}$ and define the function $\Phi_{\mathbb{W}}^{\mathbb{N}} : \mathcal{X} \longrightarrow \mathcal{B}^{\mathbb{N}}$ by

$$\Phi_{\mathbb{W}}^{\mathbb{N}}(\mathbf{x}) \quad:=\quad [\mathbf{x}_{\mathbb{W}},\ \Phi(\mathbf{x})_{\mathbb{W}},\ \Phi^2(\mathbf{x})_{\mathbb{W}},\ \Phi^3(\mathbf{x})_{\mathbb{W}}, \ldots], \qquad \text{for all } \mathbf{x} \in \mathcal{X}.$$

Then $(\mathcal{X}, \Phi)$ is posexpansive if and only if there is some finite subset $\mathbb{W} \subset \mathbb{V}$ (called a *posexpansive window*) such that the function $\Phi_{\mathbb{W}}^{\mathbb{N}}$ is an injection. If $\mathcal{Y} := \Phi_{\mathbb{W}}^{\mathbb{N}}[\mathcal{X}] \subseteq \mathcal{B}^{\mathbb{N}}$ and $\sigma : \mathcal{B}^{\mathbb{N}} \longrightarrow \mathcal{B}^{\mathbb{N}}$ is the shift map, then $\sigma(\mathcal{Y}) \subseteq \mathcal{Y}$, and $\Phi_{\mathbb{W}}^{\mathbb{N}}$ is a topological conjugacy from $(\mathcal{X}, \Phi)$ to the system $(\mathcal{Y}, \sigma)$.

For any $\mathbb{W} \subset \mathbb{V}$ and $T \in \mathbb{N}$, define $\Phi_{\mathbb{W}}^{[0...T]} : \mathcal{X} \longrightarrow \mathcal{B}^{[0...T]}$ by

$$\Phi_{\mathbb{W}}^{[0...T]}(\mathbf{x}) \quad:=\quad [\mathbf{x}_{\mathbb{W}},\ \Phi(\mathbf{x})_{\mathbb{W}},\ \Phi^2(\mathbf{x})_{\mathbb{W}}, \ldots, \Phi^T(\mathbf{x})_{\mathbb{W}}], \qquad \text{for all } \mathbf{x} \in \mathcal{X}.$$

Let $\mathbb{W}^T := \left\{\mathsf{v} \in \mathbb{V}\ ;\ \forall\ \mathbf{x}, \mathbf{x}' \in \mathcal{X},\ \left(\Phi_{\mathbb{W}}^{[0...T]}(\mathbf{x}) = \Phi_{\mathbb{W}}^{[0...T]}(\mathbf{x}')\right) \Longrightarrow \left(x_{\mathsf{v}} = x'_{\mathsf{v}}\right)\right\}$. Thus,

$$\text{for all } \mathbf{x}, \mathbf{x}' \in \mathcal{X}, \qquad \left(\Phi_{\mathbb{W}}^{[0...T]}(\mathbf{x}) = \Phi_{\mathbb{W}}^{[0...T]}(\mathbf{x}')\right) \quad\Longrightarrow\quad \left(\mathbf{x}_{\mathbb{W}^T} = \mathbf{x}'_{\mathbb{W}^T}\right). \tag{3}$$

Then we have $\mathbb{W} = \mathbb{W}^0 \subseteq \mathbb{W}^1 \subseteq \mathbb{W}^2 \subseteq \mathbb{W}^3 \subseteq \cdots$; the sequence $\{\mathbb{W}_0^t\}_{t=0}^{\infty}$ is called the $(\mathcal{X}, \Phi)$-*panorama* of $\mathbb{W}$. Clearly, $\mathbb{W}$ is a posexpansive window for Φ if and only if

$$\bigcup_{t=0}^{\infty} \mathbb{W}^t \quad=\quad \mathbb{V}. \tag{4}$$



(A) **(B)**

Figure 2: Example 2.2(b).

Shereshevsky has shown that multidimensional cellular automata can never be posexpansive. To be precise, he showed: if $(\mathbb{G},\cdot)$ is any group with growth dimension $D \geq 2$ (e.g. $\mathbb{G} = \mathbb{Z}^D$), and $\mathcal{X} \subset \mathcal{A}^{\mathbb{G}}$ is a subshift with positive topological entropy, and $\Phi : \mathcal{A}^{\mathbb{G}} \longrightarrow \mathcal{A}^{\mathbb{G}}$ is an $\mathcal{X}$-preserving cellular automaton, then the system $(\mathcal{X},\Phi)$ is not posexpansive; see [She93, Corollary 2] or [She96, Theorem 1.1]. The special case when $\mathbb{G} = \mathbb{Z}^D$ and $\mathcal{X} = \mathcal{A}^{\mathbb{Z}^D}$ was later reproved in [FMM98, Theorem 4.4]. In this section, we will generalize this result to any symbolic dynamical system satisfying some mild symmetry and mixing conditions.

Let $(\mathbb{V}, \bullet\!\!\!\rightarrow)$ be a digraph. A injection $\tau : \mathbb{V} \longrightarrow \mathbb{V}$ is a ***subisometry*** if, for all $\mathsf{v},\mathsf{w} \in \mathbb{V}$, $(\mathsf{v} \bullet\!\!\!\rightarrow \mathsf{w}) \iff (\tau(\mathsf{v}) \bullet\!\!\!\rightarrow \tau(\mathsf{w}))$. Thus, for all $\mathsf{v} \in \mathbb{V}$ and $r > 0$, we have $\tau[\mathbb{B}(\mathsf{v},r)] \subseteq \mathbb{B}[\tau(\mathsf{v}),r]$ (with equality if $\tau : \mathbb{V} \longrightarrow \mathbb{V}$ is surjective). The map τ induces a surjection $\tau_* : \mathcal{A}^{\mathbb{V}} \longrightarrow \mathcal{A}^{\mathbb{V}}$ defined by $\tau_*(\mathbf{a}) := \mathbf{a}'$ where $a'_{\mathsf{v}} := a_{\tau(\mathsf{v})}$ for all $\mathsf{v} \in \mathbb{V}$. Let $\mathcal{X} \subseteq \mathcal{A}^{\mathbb{V}}$ be a pattern space; if τ is a subisometry and $\tau_*(\mathcal{X}) = \mathcal{X}$, then we say τ is a ***subsymmetry*** of $\mathcal{X}$.

Example 2.1. Let $\mathbb{V} = \mathbb{Z}^D \times \mathbb{N}^E$ for some $D, E \geq 0$ (or some other finitely generated monoid), with the Cayley digraph structure induced by some finite generating set. Fix $\mathsf{w} \in \mathbb{V}$, and define the ***shift map*** $\tau^{\mathsf{w}} : \mathbb{V} \longrightarrow \mathbb{V}$ by $\tau^{\mathsf{w}}(\mathsf{v}) := \mathsf{v} + \mathsf{w}$; then τ is a subisometry of the Cayley digraph. If $\mathcal{X} \subseteq \mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E}$ is a subshift, then τ is a subsymmetry of $\mathcal{X}$.

Note. Subsymmetries of $\mathcal{X}$ are not necessarily injective. For example, the unilateral shift on $\mathcal{A}^{\mathbb{N}}$ is a subsymmetry, but it is many-to-one. $\diamondsuit$

Let $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ be a symbolic dynamical system. A ***subsymmetry*** of $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ is a subisometry $\tau : \mathbb{V} \longrightarrow \mathbb{V}$ such that $\tau_*[\mathcal{X}] = \mathcal{X}$ and $\tau_* \circ \Phi = \Phi \circ \tau_*$.

Example 2.2. (a) Let $\mathbb{V} = \mathbb{Z}^D \times \mathbb{N}^E$ (or any other finitely generated group), let $\mathcal{X} \subseteq \mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E}$ be a subshift, and let $\Phi : \mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E} \longrightarrow \mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E}$ be an $\mathcal{X}$-preserving cellular automaton. Then any $(\mathbb{Z}^D \times \mathbb{N}^E)$-shift is a subsymmetry of the system $(\mathcal{A}^{\mathbb{Z}^D \times \mathbb{N}^E}, \mathcal{X}, \Phi)$.

(b) Let $\psi : \mathcal{A} \times \mathcal{A} \longrightarrow \mathcal{A}$ be a binary operator (e.g. a group operator). Let $(\mathcal{A}^{\mathbb{V}}, \Phi)$ be an arbitrary symbolic dynamical system (perhaps with no symmetries), such as the one in Figure 2(A). Define $\widetilde{\mathbb{V}} := \mathbb{V} \times \mathbb{Z}$, and identify $\mathcal{A}^{\widetilde{\mathbb{V}}}$ with $(\mathcal{A}^{\mathbb{V}})^{\mathbb{Z}}$ in the obvious way; a generic element of $\mathcal{A}^{\widetilde{\mathbb{V}}}$ could be indicated as $\widetilde{\mathbf{a}} := [\mathbf{a}^n]_{n \in \mathbb{Z}}$, where $\mathbf{a}^n \in \mathcal{A}^{\mathbb{V}}$ for all $n \in \mathbb{Z}$. Let

$\sigma : (\mathcal{A}^{\mathbb{V}})^{\mathbb{Z}} \longrightarrow (\mathcal{A}^{\mathbb{V}})^{\mathbb{Z}}$ be the shift map; then σ is a subsymmetry of $\mathcal{A}^{\widetilde{\mathbb{V}}}$. Define $\widetilde{\Phi} : \mathcal{A}^{\widetilde{\mathbb{V}}} \longrightarrow \mathcal{A}^{\widetilde{\mathbb{V}}}$ by $\widetilde{\Phi}[\widetilde{\mathbf{a}}] = \widetilde{\mathbf{b}}$, where $b^n_{\mathsf{v}} = \psi[\Phi(\mathbf{a}^n)_{\mathsf{v}},\ a^{n+1}_{\mathsf{v}}]$ for all $n \in \mathbb{Z}$ and $\mathsf{v} \in \mathbb{V}$; this yields the connection network in Figure 2(B). Then σ is a subsymmetry of $(\mathcal{A}^{\widetilde{\mathbb{V}}}, \widetilde{\Phi})$. $\diamondsuit$

We say that the pattern space $\mathcal{X}$ has *weak independence* if there is some constant $\epsilon > 0$ such that, for any disjoint balls $\mathbb{B}_1, \ldots, \mathbb{B}_N \subset \mathbb{V}$,

$$\log_2 |\mathcal{X}_{\mathbb{B}_1 \sqcup \cdots \sqcup \mathbb{B}_N}| \quad \geq \quad \epsilon \sum_{n=1}^{N} \log_2 |\mathcal{X}_{\mathbb{B}_n}| \,. \tag{5}$$

This can be seen as a kind of 'topological mixing' condition —it means that the information contained in balls $\mathbb{B}_1, \ldots, \mathbb{B}_{N-1}$ has limited power to predict the contents of ball $\mathbb{B}_N$.

Example 2.3. For all $\mathsf{v} \in \mathcal{A}$, let $\mathcal{A}_{\mathsf{v}} \subset \mathcal{A}$ be a subset of cardinality at least 2. Let $\mathcal{X} := \prod_{\mathsf{v} \in \mathbb{V}} \mathcal{A}_{\mathsf{v}} \subset \mathcal{A}^{\mathbb{V}}$; then $\underline{h}(\mathcal{X}) \geq 1$, and $\mathcal{X}$ has weak independence.
In particular, the space $\mathcal{X} = \mathcal{A}^{\mathbb{V}}$ itself satisfies weak independence. $\diamondsuit$

For any $\mathsf{v} \sim \mathsf{w} \in \mathbb{V}$, let $d(\mathsf{v}, \mathsf{w})$ be the length of the shortest *undirected* path from v to w; then d is a metric on each connected component of $\mathbb{V}$. (If $\mathsf{v} \not\sim \mathsf{w}$, let $d(\mathsf{v}, \mathsf{w}) := \infty$). For any $\mathsf{v} \in \mathbb{V}$, let $\mathrm{speed}(\mathsf{v}, \tau) \ := \ \lim_{n\to\infty} \dfrac{d[\mathsf{v}, \tau^n(\mathsf{v})]}{n}$.

Lemma 2.4 **(a)** *For any* $\mathsf{v}, \mathsf{w} \in \mathbb{V}$, *we have* $d[\tau(\mathsf{v}), \tau(\mathsf{w})] \leq d(\mathsf{v}, \mathsf{w})$ (with equality if $\tau : \mathbb{V} \longrightarrow \mathbb{V}$ is surjective).

(b) *For any* $\mathsf{v} \in \mathbb{V}$, *we have* $\mathrm{speed}(\mathsf{v}, \tau) = \inf_{n \in \mathbb{N}} \dfrac{d[\mathsf{v}, \tau^n(\mathsf{v})]}{n}$.

(c) *For all* $\mathsf{v}, \mathsf{w} \in \mathbb{V}$, *if* $\mathsf{v} \sim \mathsf{w}$, *then* $\mathrm{speed}(\mathsf{v}, \tau) = \mathrm{speed}(\mathsf{w}, \tau)$.

Proof. (a) Let $(\mathsf{v}_0, \mathsf{v}_1, \mathsf{v}_2, \ldots, \mathsf{v}_N)$ be a minimal undirected path from v to w (i.e. $\mathsf{v}_0 = \mathsf{v}$, $\mathsf{v}_N = \mathsf{w}$, and either $\mathsf{v}_{n-1} \bullet\!\!\rightarrow \mathsf{v}_n$ or $\mathsf{v}_n \bullet\!\!\rightarrow \mathsf{v}_{n-1}$ for all $n \in [1...N]$). Then $(\tau(\mathsf{v}_0), \tau(\mathsf{v}_1), \ldots, \tau(\mathsf{v}_N))$ is an undirected path of length N from $\tau(\mathsf{v})$ to $\tau(\mathsf{w})$. (However, there may exist shorter paths from $\tau(\mathsf{v})$ to $\tau(\mathsf{w})$ which do not arise as τ-images of paths from v to w).

(b) is because the sequence $\{d[\mathsf{v}, \tau^n(\mathsf{v})]\}_{n=1}^{\infty}$ is subadditive:

$$d[\mathsf{v}, \tau^{n+m}(\mathsf{v})] \quad \underset{(\triangle)}{\leq} \quad d[\mathsf{v}, \tau^n(\mathsf{v})] + d[\tau^n(\mathsf{v}), \tau^{n+m}(\mathsf{v})] \quad \underset{(@)}{\leq} \quad d[\mathsf{v}, \tau^n(\mathsf{v})] + d[\mathsf{v}, \tau^m(\mathsf{v})].$$

Here $(\triangle)$ is the triangle inequality, and (@) is by part (a).

To see (c), let $r := d(\mathsf{v}, \mathsf{w})$ (finite because $\mathsf{v} \sim \mathsf{w}$). Then for any $n \in \mathbb{N}$,

$$\begin{aligned} d[\mathsf{v}, \tau^n(\mathsf{v})] \quad & \underset{(\triangle)}{\leq} \quad d[\mathsf{v}, \mathsf{w}] + d[\mathsf{w}, \tau^n(\mathsf{w})] + d[\tau^n(\mathsf{w}), \tau^n(\mathsf{v})] \\ & \underset{(@)}{\leq} \quad d[\mathsf{v}, \mathsf{w}] + d[\mathsf{w}, \tau^n(\mathsf{w})] + d[\mathsf{w}, \mathsf{v}] \quad \underset{(\dagger)}{=} \quad d[\mathsf{w}, \tau^n(\mathsf{w})] + 2r. \end{aligned}$$

Thus, $\mathrm{speed}(\mathsf{v}, \tau) \ := \ \lim_{n\to\infty} \dfrac{d[\mathsf{v}, \tau^n(\mathsf{v})]}{n} \ \leq \ \lim_{n\to\infty} \dfrac{d[\mathsf{w}, \tau^n(\mathsf{w})] + 2r}{n} \ = \ \mathrm{speed}(\mathsf{w}, \tau).$



Figure 3: Example 2.5(b).

Here, (@) is by part (a), and (†) is because $d[\mathsf{w}, \mathsf{v}] = d[\mathsf{v}, \mathsf{w}]$ (because the definition of d is symmetric) and $d[\mathsf{v}, \mathsf{w}] = r$. A symmetric argument yields $\mathrm{speed}(\mathsf{w}, \tau) \leq \mathrm{speed}(\mathsf{v}, \tau)$. Thus, $\mathrm{speed}(\mathsf{v}, \tau) = \mathrm{speed}(\mathsf{w}, \tau)$. □

Lemma 2.4(b) says the limit defining $\mathrm{speed}(\mathsf{v}, \tau)$ exists for all $\mathsf{v} \in \mathbb{V}$. We say that τ is a *moving subsymmetry* if $\mathrm{speed}(\mathsf{v}, \tau) > 0$ for all $\mathsf{v} \in \mathbb{V}$. [Lemma 2.4(c) implies that it suffices to require $\mathrm{speed}(\mathsf{v}, \tau) > 0$ for at least one v in each connected component of $(\mathbb{V}, \bullet\!\!\to)$.]

Example 2.5. (a) Let $\mathbb{V} = \mathbb{Z}^D$ with the Cayley digraph structure induced by the standard generating set $\{(\pm 1, 0, 0, \ldots, 0),\ (0, \pm 1, 0, \ldots, 0), \ldots, (0, \ldots, 0, \pm 1)\}$. If $\mathsf{z} = (z_1, \ldots, z_D) \in \mathbb{Z}^D$, and $\mathsf{o} = (0, \ldots, 0)$, then $d(\mathsf{o}, \mathsf{z}) = |z_1| + \cdots + |z_D|$. Let $\sigma^{\mathsf{z}} : \mathbb{Z}^D \longrightarrow \mathbb{Z}^D$ be the shift map. Then $\mathrm{speed}(\sigma^{\mathsf{z}}, \mathsf{v}) = d(\mathsf{o}, \mathsf{z})$ for all $\mathsf{v} \in \mathbb{Z}^D$. Thus, all nontrivial shifts are moving symmetries of $\mathcal{A}^{\mathbb{V}}$.

(b) Let $\mathbb{V} = \mathbb{Z} \times \mathbb{N}$, with the digraph structure shown in Figure 3. Here, for any $(z, n) \in \mathbb{V}$, we have $(z, n) \bullet\!\!\to (z, n')$ whenever $n' = n \pm 1$, and we also have $(z, n) \bullet\!\!\to (z', n)$ whenever $z' = z + 2^n$. Thus: $\cdots \bullet\!\!\to (-1, 0) \bullet\!\!\to (0, 0) \bullet\!\!\to (1, 0) \bullet\!\!\to (2, 0) \bullet\!\!\to (3, 0) \bullet\!\!\to \cdots$, and $\cdots \bullet\!\!\to (-2, 1) \bullet\!\!\to (0, 1) \bullet\!\!\to (2, 1) \bullet\!\!\to (4, 1) \bullet\!\!\to (6, 1) \bullet\!\!\to \cdots$, and $\cdots \bullet\!\!\to (-4, 2) \bullet\!\!\to (0, 2) \bullet\!\!\to (4, 2) \bullet\!\!\to (8, 2) \bullet\!\!\to (12, 2) \bullet\!\!\to \cdots$, etc. Define subisometry $\tau : \mathbb{V} \longrightarrow \mathbb{V}$ by $\tau(z, n) = (z + 1, n)$. Then $\mathrm{speed}(\tau, \mathsf{v}) = 0$, for all $\mathsf{v} \in \mathbb{V}$, because for any $k \in \mathbb{N}$, there is a path from v to $\tau^{(2^k)}(\mathsf{v})$ of length at most $2k + 1$. Thus, τ is *not* a moving subsymmetry. ♢

In a digraph $(\mathbb{V}, \bullet\!\!\to)$, a vertex $\mathsf{v} \in \mathbb{V}$ has *superlinear connectivity* if $\liminf_{r \to \infty} \dfrac{|\mathbb{B}(\mathsf{v}, r)|}{r} = \infty$.

Example 2.6. (a) If $\underline{\dim}_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\to) > 1$, then v has superlinear connectivity. (For example, if $\mathbb{V}$ is a Cayley digraph of a group with growth dimension $D \geq 2$, then every vertex has superlinear connectivity.)

(b) If $\mathsf{v} \rightsquigarrow \mathsf{w}$ and v has superlinear connectivity, then w has superlinear connectivity. ♢

The main result of this section is the following:

Theorem 2.7 *Let $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ be a symbolic dynamical system with a moving subsymmetry. If $\mathcal{X}$ has weak independence, and there exists some $\mathsf{v} \in \mathbb{V}$ with superlinear connectivity such that $\overline{h}_{\mathsf{v}}(\mathcal{X}) > 0$, then the system $(\mathcal{X}, \Phi)$ is not posexpansive.*

Before proving Theorem 2.7, we give two concrete corollaries.

Corollary 2.8 *Let* $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ *be a symbolic dynamical system with a moving subsymmetry, such that* $\mathcal{X}$ *has weak independence. Suppose that either*

> **(a)** $\overline{h}(\mathcal{X}) > 0$ *and* $\underline{\dim}(\mathbb{V}, \bullet\!\!\!\rightarrow) > 1$*; or*
>
> **(b)** $(\mathbb{V}, \bullet\!\!\!\rightarrow)$ *is dimensionally regular,* $\underline{h}(\mathcal{X}) > 0$*, and* $\overline{\dim}(\mathbb{V}, \bullet\!\!\!\rightarrow) > 1$.

Then the system $(\mathcal{X}, \Phi)$ *is not posexpansive.*

Proof. Recall: if $\underline{\dim}_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\!\rightarrow) > 1$, then v has superlinear connectivity.

> (a) There exists $\mathsf{v} \in \mathbb{V}$ with $\overline{h}_{\mathsf{v}}(\mathcal{X}) > 0$, because $\overline{h}(\mathcal{X}) > 0$. But v also has superlinear connectivity, because $\underline{\dim}_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\!\rightarrow) \geq \underline{\dim}(\mathbb{V}, \bullet\!\!\!\rightarrow) > 1$. Now apply Theorem 2.7.
>
> (b) There exists $\mathsf{v} \in \mathbb{V}$ with $\overline{\dim}_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\!\rightarrow) > 1$, because $\overline{\dim}(\mathbb{V}, \bullet\!\!\!\rightarrow) > 1$. Thus, $\underline{\dim}_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\!\rightarrow) > 1$ also, because $(\mathbb{V}, \bullet\!\!\!\rightarrow)$ is dimensionally regular. Thus, v has superlinear connectivity. Also, $\overline{h}_{\mathsf{v}}(\mathcal{X}) > \underline{h}_{\mathsf{v}}(\mathcal{X}) \geq \underline{h}(\mathcal{X}) > 0$. Now apply Theorem 2.7. □

Corollary 2.9 *Let* $\Phi : \mathcal{A}^{\mathbb{V}} \longrightarrow \mathcal{A}^{\mathbb{V}}$ *be a continuous self-map with a moving subsymmetry. If* $\underline{\dim}(\mathbb{V}, \bullet\!\!\!\rightarrow) > 1$*, then the system* $(\mathcal{A}^{\mathbb{V}}, \Phi)$ *is not posexpansive.*

Proof. If $\mathcal{X} = \mathcal{A}^{\mathbb{V}}$, then clearly $\mathcal{X}$ has weak independence, and $\overline{h}(\mathcal{X}) = \log_2|\mathcal{A}| > 0$. Now apply Corollary 2.8(a). □

The proof of Theorem 2.7 consists of two lemmas concerning the 'entropy' of a pattern space relative to a subsymmetry. Let $\mathcal{X} \subseteq \mathcal{A}^{\mathbb{V}}$ be a pattern space and let $\tau : \mathbb{V} \longrightarrow \mathbb{V}$ be a subsymmetry of $\mathcal{X}$. For any finite $\mathbb{F} \subset \mathbb{V}$, we define

$$\overline{h}(\mathcal{X}, \tau, \mathbb{F}) \quad := \quad \limsup_{N\to\infty} \frac{1}{N} \log_2 \left|\mathcal{X}_{\mathbb{F}(N)}\right|, \quad \text{where} \quad \mathbb{F}(N) \ := \ \bigcup_{n=0}^{N} \tau^n(\mathbb{F}) \ \subseteq \ \mathbb{V}. \tag{6}$$

We then define the ***upper*** τ***-entropy*** of $\mathcal{X}$ by

$$\overline{h}(\mathcal{X}, \tau) \quad := \quad \sup_{\substack{\mathbb{F}\subset\mathbb{V} \\ \text{finite}}} \overline{h}(\mathcal{X}, \tau, \mathbb{F}). \tag{7}$$

Lemma 2.10 *Let* $(\mathbb{V}, \bullet\!\!\!\rightarrow)$ *be a digraph, and let* $\mathcal{X} \subseteq \mathcal{A}^{\mathbb{V}}$ *be a pattern space with weak independence. Suppose there exists* $\mathsf{v} \in \mathbb{V}$ *with superlinear connectivity and* $\overline{h}_{\mathsf{v}}(\mathcal{X}) > 0$*. If* $\tau : \mathbb{V} \longrightarrow \mathbb{V}$ *is any moving subsymmetry of* $\mathcal{X}$*, then* $\overline{h}(\mathcal{X}, \tau) = \infty$.

Proof. Let $\epsilon > 0$ be as in equation (5). Let $S := \text{speed}(\tau, \mathsf{v}) > 0$.

Claim 1: *For any* $r > 0$*, we have* $\overline{h}[\mathcal{X}, \tau, \mathbb{B}(\mathsf{v}, r)] \ \geq \ \dfrac{S\epsilon}{4r} \ \log_2\left|\mathcal{X}_{\mathbb{B}(\mathsf{v},r)}\right|$.

Proof. Let $m := \lceil 2r/S \rceil$, then the points $\{\mathsf{v},\ \tau^m(\mathsf{v}),\ \tau^{2m}(\mathsf{v}),\ \tau^{3m}(\mathsf{v}), \ldots\}$ are all at least $2r$-separated, by Lemma 2.4(b). Thus, the balls $\{\mathbb{B}(\mathsf{v}, r),\ \mathbb{B}(\tau^m(\mathsf{v}), r),\ \mathbb{B}(\tau^{2m}(\mathsf{v}), r), \ldots\}$ are all disjoint. Let $\mathbb{F} := \mathbb{B}(\mathsf{v}, r)$; then for any $n \in \mathbb{N}$, we have $\tau^{nm}[\mathbb{F}] \subseteq \mathbb{B}(\tau^{nm}(\mathsf{v}), r)$ (because τ is a subisometry of $\mathbb{V}$). Thus, the sets $\{\mathbb{F},\ \tau^m(\mathbb{F}),\ \tau^{2m}(\mathbb{F}), \ldots\}$ are disjoint.

For any $N \in \mathbb{N}$, let $\mathbb{F}(Nm) := \bigcup_{k=0}^{Nm} \tau^k(\mathbb{F})$; then $\mathbb{F}(Nm) \supseteq \bigsqcup_{n=0}^{N} \tau^{nm}(\mathbb{F})$. Thus

$$\log_2 \left|\mathcal{X}_{\mathbb{F}(Nm)}\right| \quad \underset{(*)}{\geq} \quad \epsilon \cdot \sum_{n=1}^{N} \log_2 \left|\mathcal{X}_{\tau^{nm}(\mathbb{F})}\right| \quad \underset{(\dagger)}{=} \quad \epsilon N \log_2 |\mathcal{X}_{\mathbb{F}}|, \tag{8}$$

where $(*)$ is by equation (5), and $(\dagger)$ is because τ is a subsymmetry of $\mathcal{X}$ (so $|\mathcal{X}_{\mathbb{F}}| = \left|\mathcal{X}_{\tau^k(\mathbb{F})}\right|$ for all $k \in \mathbb{Z}$). Combining equations (6) and (8), we get

$$\begin{aligned}
\overline{h}(\mathcal{X}, \tau, \mathbb{F}) \quad & \underset{(6)}{:=} \quad \limsup_{N\to\infty} \frac{1}{N} \log_2 \left|\mathcal{X}_{\mathbb{F}(N)}\right| \quad \geq \quad \limsup_{N\to\infty} \frac{1}{Nm} \log_2 \left|\mathcal{X}_{\mathbb{F}(Nm)}\right| \\
& \underset{(8)}{\geq} \quad \lim_{N\to\infty} \frac{\epsilon\, N}{N\, m} \log_2 |\mathcal{X}_{\mathbb{F}}| \quad = \quad \frac{\epsilon}{m} \cdot \log_2 |\mathcal{X}_{\mathbb{F}}| \quad \underset{(*)}{\geq} \quad \frac{S\epsilon}{4r} \cdot \log_2 |\mathcal{X}_{\mathbb{F}}|,
\end{aligned}$$

as desired. Here, $(*)$ is because $m := \lceil 2r/S \rceil \leq 4r/S$. ◇ `Claim 1`

It follows from defining equation (7) that

$$\begin{aligned}
\overline{h}[\mathcal{X}, \tau] \quad & \underset{(7)}{\geq} \quad \sup_{r\in\mathbb{N}} \ \overline{h}[\mathcal{X}, \tau, \mathbb{B}(\mathsf{v}, r)] \quad \geq \quad \limsup_{r\to\infty} \ \overline{h}[\mathcal{X}, \tau, \mathbb{B}(\mathsf{v}, r)] \quad \underset{(\dagger)}{\geq} \quad \limsup_{r\to\infty} \ \frac{S\epsilon}{4r} \log_2 \left|\mathcal{X}_{\mathbb{B}(\mathsf{v},r)}\right| \\
& = \quad \limsup_{r\to\infty} \left(\frac{S\epsilon}{4} \cdot \frac{\log_2 \left|\mathcal{X}_{\mathbb{B}(\mathsf{v},r)}\right|}{|\mathbb{B}(\mathsf{v}, r)|} \cdot \frac{|\mathbb{B}(\mathsf{v}, r)|}{r} \right) \\
& \geq \quad \frac{S\epsilon}{4} \cdot \left(\limsup_{r\to\infty} \ \frac{\log_2 \left|\mathcal{X}_{\mathbb{B}(\mathsf{v},r)}\right|}{|\mathbb{B}(\mathsf{v}, r)|} \right) \cdot \left(\liminf_{r\to\infty} \ \frac{|\mathbb{B}(\mathsf{v}, r)|}{r} \right) \\
& \underset{(2)}{=} \quad \left(\frac{S\epsilon\, \overline{h}_{\mathsf{v}}(\mathcal{X})}{4} \right) \ \liminf_{r\to\infty} \ \frac{|\mathbb{B}(\mathsf{v}, r)|}{r} \quad \underset{(*)}{=} \quad \infty,
\end{aligned}$$

as desired. Here, $(\dagger)$ is by Claim 1, and $(*)$ is because $\overline{h}_{\mathsf{v}}(\mathcal{X}) > 0$ and $(\mathbb{V}, \bullet\!\!\!\rightarrow)$ has superlinear connectivity at v. □

Lemma 2.11 *Let $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ symbolic dynamical system with a subsymmetry $\tau : \mathbb{V} \longrightarrow \mathbb{V}$. If $\overline{h}(\mathcal{X}, \tau) = \infty$, then $(\mathcal{X}, \Phi)$ is not posexpansive.*

Proof. (by contradiction) Suppose $(\mathcal{X}, \Phi)$ is posexpansive. Let $\mathbb{W}_0 \subseteq \mathbb{V}$ be a posexpansive window, with panorama $\{\mathbb{W}_0^t\}_{t=0}^{\infty}$. For any $n \in \mathbb{N}$, let $\mathbb{W}_n := \tau^n(\mathbb{W}_0)$, and for all $t \in \mathbb{N}$, let $\mathbb{W}_n^t := \tau^n(\mathbb{W}_0^t)$.

Claim 1: *For all $n \in \mathbb{N}$, $\{\mathbb{W}_n^t\}_{t=0}^{\infty}$ is the panorama of $\mathbb{W}_n$.*

Proof. For any $\mathbf{x}, \mathbf{x}' \in \mathcal{X}$, and any $T \in \mathbb{N}$, we have

$$\begin{aligned}
\left(\Phi^{[0...T]}_{\mathbb{W}_n}(\mathbf{x}) = \Phi^{[0...T]}_{\mathbb{W}_n}(\mathbf{x}')\right) \quad &\Longleftrightarrow \quad \left(\Phi^t(\mathbf{x})_{\mathbb{W}_n} = \Phi^t(\mathbf{x}')_{\mathbb{W}_n},\ \ \forall\ t \in [0...T]\right) \\
\Longleftrightarrow \quad &\left(\tau^n \circ \Phi^t(\mathbf{x})_{\mathbb{W}_0} = \tau^n \circ \Phi^t(\mathbf{x}')_{\mathbb{W}_0}, \forall\ t \in [0...T]\right) \\
\underset{(\dagger)}{\Longleftrightarrow} \quad &\left(\Phi^t \circ \tau^n(\mathbf{x})_{\mathbb{W}_0} = \Phi^t \circ \tau^n(\mathbf{x}')_{\mathbb{W}_0}, \forall\ t \in [0...T]\right) \\
\Longleftrightarrow \quad &\left(\Phi^{[0...T]}_{\mathbb{W}_0}[\tau^n(\mathbf{x})] = \Phi^{[0...T]}_{\mathbb{W}_0}[\tau^n(\mathbf{x}')]\right) \\
\underset{(*)}{\Longrightarrow} \quad &\left(\tau^n(\mathbf{x})_{\mathbb{W}_0^T} = \tau^n(\mathbf{x}')_{\mathbb{W}_0^T}\right) \quad \Longleftrightarrow \quad \left(\mathbf{x}_{\mathbb{W}_n^T} = \mathbf{x}'_{\mathbb{W}_n^T}\right),
\end{aligned}$$

as desired. Here $(*)$ is by statement (3), because $\{\mathbb{W}_0^t\}_{t=0}^{\infty}$ is the panorama of $\mathbb{W}_0$, and $(\dagger)$ is because $\Phi \circ \tau^n = \tau^n \circ \Phi$, because τ is a subsymmetry of $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$. $\diamond$ `Claim 1`

Claim 2: (a) *There exists $T \in \mathbb{N}$ such that $\mathbb{W}_1 \subseteq \mathbb{W}_0^T$.*

(b) *For all $t \in \mathbb{N}$, we have $\mathbb{W}_1^t \subseteq \mathbb{W}_0^{T+t}$.*

(c) *For all $n \in \mathbb{N}$, we have $\mathbb{W}_n \subseteq \mathbb{W}_0^{nT}$.*

(d) *For all $n, t \in \mathbb{N}$, we have $\mathbb{W}_n^t \subseteq \mathbb{W}_0^{nT+t}$.*

Proof. (a) follows from equation (4). To see (b), let $\mathbf{x}, \mathbf{x}' \in \mathcal{X}$. Then

$$\begin{aligned}
\left(\Phi^{[0...T+t]}_{\mathbb{W}_0}(\mathbf{x}) = \Phi^{[0...T+t]}_{\mathbb{W}_0}(\mathbf{x}')\right) \quad \Longleftrightarrow \quad &\left(\Phi^{[s...T+s]}_{\mathbb{W}_0}(\mathbf{x}) = \Phi^{[s...T+s]}_{\mathbb{W}_0}(\mathbf{x}') \text{ for all } s \in [0...t]\right) \\
\underset{(\dagger)}{\Longleftrightarrow} \quad &\left(\Phi^{[0...T]}_{\mathbb{W}_0}[\Phi^s(\mathbf{x})] = \Phi^{[0...T]}_{\mathbb{W}_0}[\Phi^s(\mathbf{x}')] \text{ for all } s \in [0...t]\right) \\
\underset{(*)}{\Longrightarrow} \quad &\left(\Phi^s(\mathbf{x})_{\mathbb{W}_0^T} = \Phi^s(\mathbf{x}')_{\mathbb{W}_0^T} \text{ for all } s \in [0...t]\right) \\
\underset{(@)}{\Longrightarrow} \quad &\left(\Phi^s(\mathbf{x})_{\mathbb{W}_1} = \Phi^s(\mathbf{x}')_{\mathbb{W}_1} \text{ for all } s \in [0...t]\right) \\
\underset{(\diamond)}{\Longleftrightarrow} \quad &\left(\Phi^{[0...t]}_{\mathbb{W}_1}(\mathbf{x}) = \Phi^{[0...t]}_{\mathbb{W}_1}(\mathbf{x}')\right) \underset{(\ddagger)}{\Longrightarrow} \left(\mathbf{x}_{\mathbb{W}_1^t} = \mathbf{x}'_{\mathbb{W}_1^t}\right).
\end{aligned}$$

Thus, $\mathbb{W}_1^t \subseteq \mathbb{W}_0^{T+t}$, as desired. Here, $(*)$ is by statement (3), because $\{\mathbb{W}_0^t\}_{t=0}^{\infty}$ is the panorama of $\mathbb{W}_0$. $(\dagger)$ is because $\Phi^{[s...T+s]}_{\mathbb{W}_0}(\mathbf{x}) = \Phi^{[0...T]}_{\mathbb{W}_0}[\Phi^s(\mathbf{x})]$ (and likewise for $\mathbf{x}'$). $(@)$ is by part (a). Finally, $(\diamond)$ is by definition of $\Phi^{[0...t]}_{\mathbb{W}_1}(\mathbf{x})$, and $(\ddagger)$ is by statement (3) and Claim 1.

(c) (by induction on n) The case $(n = 1)$ is Part (a). Now suppose inductively that $\mathbb{W}_n \subseteq \mathbb{W}_0^{nT}$. Then $\mathbb{W}_{n+1} = \tau(\mathbb{W}_n) \underset{(\dagger)}{\subseteq} \tau(\mathbb{W}_0^{nT}) = \mathbb{W}_1^{nT} \underset{(*)}{\subseteq} \mathbb{W}_0^{nT+T} = \mathbb{W}_0^{(n+1)T}$. Here, $(\dagger)$ is by the induction hypothesis, and $(*)$ is by part (b).

(d) is the same argument as in part (b), but using part (c) instead of part (a). $\diamond$ `Claim 2`

Now, for any $H > 0$, we can find some finite subset $\mathbb{F} \subset \mathbb{V}$ such that $\overline{h}(\mathcal{X}, \tau, \mathbb{F}) > H$ (because $\overline{h}(\mathcal{X}, \tau) = \infty$). Equation (4) yields some t such that $\mathbb{F} \subseteq \mathbb{W}_0^t$. Thus, for any $N \in \mathbb{N}$, and all $n \in [0...N]$, we have

$$\tau^n[\mathbb{F}] \quad \subseteq \quad \tau^n[\mathbb{W}_0^t] \quad = \quad \mathbb{W}_n^t \quad \underset{(*)}{\subseteq} \quad \mathbb{W}_0^{nT+t} \quad \underset{(\dagger)}{\subseteq} \quad \mathbb{W}_0^{NT+t},$$

where $(*)$ is by Claim 2(d), and $(\dagger)$ is because $nT+t \leq NT+t$. Thus,

$$\text{if} \quad \mathbb{F}(N) \quad := \quad \bigcup_{n=0}^{N} \tau^n(\mathbb{F}), \quad \text{then} \quad \mathbb{W}_0^{NT+t} \quad \supseteq \quad \mathbb{F}(N). \tag{9}$$

Thus,

$$\log_2 \left|\Phi_{\mathbb{W}_0}^{[0\ldots NT+t]}(\mathcal{X})\right| \quad \underset{(*)}{\geq} \quad \log_2 \left|\mathcal{X}_{\mathbb{W}_0^{NT+t}}\right| \quad \underset{(\dagger)}{\geq} \quad \log_2 \left|\mathcal{X}_{\mathbb{F}(N)}\right|. \tag{10}$$

Here, $(*)$ is because statement (3) yields an injection from $\mathcal{X}_{\mathbb{W}_0^{NT+t}}$ into $\Phi_{\mathbb{W}_0}^{[0\ldots NT+t]}(\mathcal{X})$; meanwhile $(\dagger)$ is by equation (9). Let $\mathcal{B} := \mathcal{A}^{\mathbb{W}_0}$ and $\mathcal{Y} := \Phi_{\mathbb{W}_0}^{\mathbb{N}}(\mathcal{X}) \subseteq \mathcal{B}^{\mathbb{N}}$. Then

$$\begin{aligned}
\overline{h}(\mathcal{Y},\sigma) \quad &:= \quad \limsup_{M\to\infty} \frac{\log_2 |\mathcal{Y}_{[0\ldots M]}|}{M} \quad \geq \quad \limsup_{N\to\infty} \frac{\log_2 |\mathcal{Y}_{[0\ldots NT+t]}|}{NT+t} \\
&= \quad \limsup_{N\to\infty} \frac{\log_2 \left|\Phi_{\mathbb{W}_0}^{[0\ldots NT+t]}(\mathcal{X})\right|}{NT+t} \quad \underset{(\dagger)}{\geq} \quad \limsup_{N\to\infty} \frac{\log_2 \left|\mathcal{X}_{\mathbb{F}(N)}\right|}{NT+t} \\
&= \quad \limsup_{N\to\infty} \left(\frac{N}{NT+t}\right)\left(\frac{\log_2 \left|\mathcal{X}_{\mathbb{F}(N)}\right|}{N}\right) \quad \underset{(\ddagger)}{=} \quad \frac{\overline{h}(\mathcal{X},\tau,\mathbb{F})}{T} \quad > \quad \frac{H}{T}.
\end{aligned} \tag{11}$$

Here, $(\dagger)$ is by equation (10), and $(\ddagger)$ is by defining equation (6). Now, H can be made arbitrarily large, because $\overline{h}(\mathcal{X},\tau) = \infty$. Thus, letting $H\to\infty$ in equation (11), we conclude that $\overline{h}(\mathcal{Y},\sigma) = \infty$.

But clearly, $\overline{h}(\mathcal{Y},\sigma) \leq \log_2 |\mathcal{B}| = \log_2 |\mathcal{A}^{\mathbb{W}_0}| = |\mathbb{W}_0| \cdot \log_2 |\mathcal{A}| < \infty$, because $\mathcal{A}$ and $\mathbb{W}_0$ are finite. Contradiction. □

Proof of Theorem 2.7. Combine Lemmas 2.10 and 2.11. □

Remarks. (a) Observe that Lemma 2.11 is really a statement about $(\mathcal{X},\Phi)$ as an abstract Cantor dynamical system with a subsymmetry; it does not depend on any specific representation of $(\mathcal{X},\Phi)$ as a symbolic dynamical system (i.e. any specific embedding $\mathcal{X} \subset \mathcal{A}^{\mathbb{V}}$ for some $\mathcal{A}$ and $\mathbb{V}$). As such, Lemma 2.11 is an interesting result in itself.

(b) Theorem 2.7 applies even if τ and its iterates are the only symmetries of $(\mathcal{A}^{\mathbb{V}},\mathcal{X},\Phi)$. In particular, we do *not* require the symmetry group of $(\mathcal{A}^{\mathbb{V}},\mathcal{X},\Phi)$ to itself have growth dimension greater than 1.

(c) The 'weak independence' condition in Theorem 2.7 and Lemma 2.10 is probably not necessary. ◇.

3 Propagation, Sensitivity, and Equicontinuity

If $\Phi : \mathcal{A}^{\mathbb{V}} \longrightarrow \mathcal{A}^{\mathbb{V}}$ is continuous, then for all $t \in \mathbb{N}$, the function $\Phi^t : \mathcal{A}^{\mathbb{V}} \longrightarrow \mathcal{A}^{\mathbb{V}}$ is also continuous; hence we can apply Lemma 1.2 to define input neighbourhoods $\Phi^t_{\text{in}}(\mathsf{v}) \subset \mathbb{V}$ for all $\mathsf{v} \in \mathbb{V}$. The ***propagation*** of Φ at v is the function $\rho_{\mathsf{v}} : \mathbb{N} \longrightarrow \mathbb{N}$ defined by

$$\rho_{\mathsf{v}}(T) := \left|\Phi^{[0...T]}_{\text{in}}(\mathsf{v})\right|, \quad \text{where} \quad \Phi^{[0...T]}_{\text{in}}(\mathsf{v}) \ := \ \bigcup_{t=0}^{T} \Phi^t_{\text{in}}(\mathsf{v}), \quad \text{for all } T \in \mathbb{N}. \tag{12}$$

Clearly, $\rho_{\mathsf{v}}(T) \leq |\mathbb{B}(\mathsf{v}, T)|$ (because for all $t \leq T$, we have $\Phi^t_{\text{in}}(\mathsf{v}) \ \subseteq \ \mathbb{B}(\mathsf{v}, T)$). In general, this inequality may be strict.

Let $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ be a symbolic dynamical system, and let $\mathsf{v} \in \mathbb{V}$. A point $\mathbf{x} \in \mathcal{X}$ is v-***sensitive*** if there exists a sequence $\{\mathbf{x}^n\}_{n=1}^{\infty} \subset \mathcal{X}$ with $\lim_{n\to\infty} \mathbf{x}^n \ = \ \mathbf{x}$, such that

$$\text{For all } n \in \mathbb{N}, \text{ there is some } t \in \mathbb{N} \text{ with } \Phi^t(\mathbf{x}^n)_{\mathsf{v}} \neq \Phi^t(\mathbf{x})_{\mathsf{v}}. \tag{13}$$

We say that $\mathbf{x}$ is a ***sensitive point*** if it is v-sensitive for some $\mathsf{v} \in \mathbb{V}$. (If d is any compatible metric on $\mathcal{X}$, then there is some ϵ such that, for all $\mathbf{x}, \mathbf{y} \in \mathcal{X}$, $\Big(x_{\mathsf{v}} \neq y_{\mathsf{v}}\Big) \Longrightarrow \Big(d(\mathbf{x}, \mathbf{y}) > \epsilon\Big)$; thus, this definition is equivalent to the ordinary metric definition of 'sensitivity').

Proposition 3.1 *Let $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ be a symbolic dynamical system.*

(a) *Let $\mathsf{v} \in \mathbb{V}$ and suppose Φ has propagation ρ_{v} at v. Then $\Big(\rho_{\mathsf{v}}$ is unbounded$\Big) \iff \Big((\mathcal{X}, \Phi)$ has a v-sensitive point$\Big)$.*

(b) *$(\mathcal{X}, \Phi)$ has a sensitive point if and only if there exists $\mathsf{v} \in \mathbb{V}$ with unbounded propagation.*

Proof. (a) "$\Longrightarrow$" For all $r \in \mathbb{N}$, there exists $T(r) \in \mathbb{N}$ such that $\rho_{\mathsf{v}}[T(r)] > |\mathbb{B}(\mathsf{v}, r)|$, which means there is some $t(r) \in [0...T(r)]$ with $\Phi^{t(r)}_{\text{in}}(\mathsf{v}) \not\subseteq \mathbb{B}(\mathsf{v}, r)$. Let $\mathsf{w} \in \Phi^{t(r)}_{\text{in}}(\mathsf{v}) \setminus \mathbb{B}(\mathsf{v}, r)$. The local rule $\phi^{t(r)}_{\mathsf{v}} : \mathcal{A}^{\Phi^{t(r)}_{\text{in}}(\mathsf{v})} \longrightarrow \mathcal{A}$ is proper, so there exist $\mathbf{y}^r, \mathbf{z}^r \in \mathcal{X}$ such that

$$(\mathbf{a}) \quad y^r_{\mathsf{u}} \ = \ z^r_{\mathsf{u}}, \text{ for all } \mathsf{u} \in \mathbb{V} \setminus \{\mathsf{w}\}; \quad \text{but} \quad (\mathbf{b}) \quad \Phi^{t(r)}(\mathbf{y}^r)_{\mathsf{v}} \ \neq \ \Phi^{t(r)}(\mathbf{z}^r)_{\mathsf{v}}.$$

Now, $\mathbb{B}(\mathsf{v}, r) \subseteq \mathbb{V} \setminus \{\mathsf{w}\}$ by construction, so condition (a) means that

$$\mathbf{y}^r_{\mathbb{B}(\mathsf{v},r)} \quad = \quad \mathbf{z}^r_{\mathbb{B}(\mathsf{v},r)}. \tag{14}$$

Since $\mathcal{X}$ is compact, we find some increasing sequence $\{r_n\}_{n=1}^{\infty} \in \mathbb{N}$ such that the subsequence $\{\mathbf{y}^{r_n}\}_{n=1}^{\infty}$ converges in $\mathcal{X}$ to some point $\mathbf{x}$. Equation (14) implies that the subsequence $\{\mathbf{z}^{r_n}\}_{n=1}^{\infty}$ also converges to $\mathbf{x}$. But for all $n \in \mathbb{N}$, condition (b) says that $\Phi^{t(r_n)}(\mathbf{y}^{r_n})_{\mathsf{v}} \neq \Phi^{t(r_n)}(\mathbf{z}^{r_n})_{\mathsf{v}}$, which means that either (i) $\Phi^{t(r_n)}(\mathbf{y}^{r_n})_{\mathsf{v}} \neq \Phi^{t(r_n)}(\mathbf{x})_{\mathsf{v}}$ or (ii) $\Phi^{t(r_n)}(\mathbf{z}^{r_n})_{\mathsf{v}} \neq \Phi^{t(r_n)}(\mathbf{x})_{\mathsf{v}}$ (or both). In case (i), define $\mathbf{x}^n := \mathbf{y}^{r_n}$, while in case (ii), define $\mathbf{x}^n := \mathbf{z}^{r_n}$; then we obtain a sequence $\{\mathbf{x}^n\}_{n=1}^{\infty}$ converging to $\mathbf{x}$, and satisfying condition (13).

"$\Longleftarrow$" For any $R \in \mathbb{N}$, we must find some $T \in \mathbb{N}$ such that $\rho_{\mathsf{v}}(T) > R$. Let $\mathbf{x} \in \mathcal{X}$ be a v-sensitive point, and let $\{\mathbf{x}^n\}_{n=1}^{\infty} \subset \mathcal{X}$ be a sequence converging to $\mathbf{x}$ and satisfying condition (13). Now, there exist some $n \in \mathbb{N}$ such that $\mathbf{x}^n_{\mathbb{B}(\mathsf{v},R)} = \mathbf{x}_{\mathbb{B}(\mathsf{v},R)}$; but there also exists some T such that $\Phi^T(\mathbf{x}^n)_{\mathsf{v}} \neq \Phi^T(\mathbf{x})_{\mathsf{v}}$. This means that $\Phi^T_{\text{in}}(\mathsf{v}) \not\subseteq \mathbb{B}(\mathsf{v},R)$. Let $\mathsf{w}_T \in \Phi^T_{\text{in}}(\mathsf{v}) \setminus \mathbb{B}(\mathsf{v},R)$. Then there exists a directed path $\mathsf{w}_T \bullet\!\!\rightarrow \mathsf{w}_{T-1} \bullet\!\!\rightarrow \cdots \bullet\!\!\rightarrow \mathsf{w}_2 \bullet\!\!\rightarrow \mathsf{w}_1 \bullet\!\!\rightarrow \mathsf{v}$ such that $\mathsf{w}_t \in \Phi^t_{\text{in}}(\mathsf{v})$ for all $t \in [1...T]$. Furthermore, this path must have length $L > R$ (even after removing repeated entries), because $\mathsf{w}_T \notin \mathbb{B}(\mathsf{v},R)$. Thus, $\rho_{\mathsf{v}}(T) := \left|\Phi^{[0..T]}_{\text{in}}(\mathsf{v})\right| \geq |\{\mathsf{w}_T, \ldots, \mathsf{w}_2, \mathsf{w}_1\}| = L > R$, as desired. This works for any $R \in \mathbb{N}$; hence the function ρ_{v} is unbounded.

(b) follows immediately from (a). ☐

Let $\mathbb{W} \subset \mathbb{V}$ be some finite subset. We say that Φ is $\mathbb{W}$-*equicontinuous* if there exists a finite subset $\mathbb{U} \subset \mathbb{V}$ containing $\mathbb{W}$ (called the *envelope* of $\mathbb{W}$), such that:

$$\text{For all } \mathbf{x}, \mathbf{y} \in \mathcal{X}, \qquad \Big(\mathbf{y}_{\mathbb{U}} = \mathbf{x}_{\mathbb{U}}\Big) \quad \Longrightarrow \quad \Big(\Phi^t(\mathbf{y})_{\mathbb{W}} = \Phi^t(\mathbf{x})_{\mathbb{W}},\ \forall t \in \mathbb{N}\Big). \tag{15}$$

We say that Φ is *equicontinuous* if Φ is $\mathbb{W}$-equicontinuous for every finite subset $\mathbb{W} \subset \mathbb{V}$. (If d is any compatible metric on $\mathcal{X}$, then for any $\epsilon > 0$ there is some finite subset $\mathbb{W} \subset \mathbb{V}$ such that for all $\mathbf{x}, \mathbf{y} \in \mathcal{X}$, $\Big(\mathbf{x}_{\mathbb{W}} = \mathbf{y}_{\mathbb{W}}\Big) \Longrightarrow \Big(d(\mathbf{x},\mathbf{y}) < \epsilon\Big)$. Likewise, for any finite $\mathbb{U} \subset \mathbb{V}$, there is some $\delta > 0$ such that for all $\mathbf{x}, \mathbf{y} \in \mathcal{X}$, $\Big(d(\mathbf{x},\mathbf{y}) < \delta\Big) \Longrightarrow \Big(\mathbf{x}_{\mathbb{U}} = \mathbf{y}_{\mathbb{U}}\Big)$. Thus, our definition is equivalent to the ordinary metric definition of 'equicontinuity').

A topological dynamical system $(\mathcal{X}, \Phi)$ is an *odometer* if $(\mathcal{X}, \Phi)$ is an inverse limit of a sequence of finite, periodic dynamical systems. That is:

$$(\mathcal{X}, \Phi) \quad := \quad \lim \Big\{ \cdots \xrightarrow{\pi_3} (\mathcal{X}_3, \phi_3) \xrightarrow{\pi_2} (\mathcal{X}_2, \phi_2) \xrightarrow{\pi_1} (\mathcal{X}_1, \phi_1) \Big\}, \tag{16}$$

where, for all $n \in \mathbb{N}$, $\mathcal{X}_n$ is a finite set, $\phi_n : \mathcal{X}_n \longrightarrow \mathcal{X}_n$ is a cyclic permutation, and $\pi_n : (\mathcal{X}_{n+1}, \phi_{n+1}) \longrightarrow (\mathcal{X}_n, \phi_n)$ is a factor mapping.

For any $x \in \mathcal{X}$, let $\overline{\mathcal{O}}_x := \overline{\{\Phi^t(x)\ ;\ t \in \mathbb{N}\}}$ be the Φ-*orbit closure* of x; then $(\overline{\mathcal{O}}_x, \Phi)$ is itself a topological dynamical system. The system $(\mathcal{X}, \Phi)$ is an *odometer bundle* if, for every $x \in \mathcal{X}$, the system $(\overline{\mathcal{O}}_x, \Phi)$ is an odometer. Thus, $(\mathcal{X}, \Phi)$ can be decomposed into a (possibly infinite) disjoint union of (possible non-isomorphic) odometers.

For example, for all $n \in \mathbb{N}$, let $\mathcal{X}_n$ be a finite set, and let $\phi_n : \mathcal{X}_n \longrightarrow \mathcal{X}_n$ be a permutation (possibly with multiple disjoint orbits). Suppose $(\mathcal{X}, \Phi)$ arises as the inverse limit (16); then $(\mathcal{X}, \Phi)$ is an odometer bundle.

Proposition 3.2 *For all* $\mathsf{v} \in \mathbb{V}$*, let* $\rho_{\mathsf{v}} : \mathbb{N} \longrightarrow \mathbb{N}$ *be the propagation of* Φ *at* v.

(a) *Let* $\mathbb{W} \subset \mathbb{V}$ *be a finite subset. If* ρ_{w} *is bounded for all* $\mathsf{w} \in \mathbb{W}$*, then* Φ *is* $\mathbb{W}$*-equicontinuous.*

(b) $\Big(\rho_{\mathsf{v}}$ *is bounded for all* $\mathsf{v} \in \mathbb{V}\Big) \iff \Big(\Phi$ *is equicontinuous*$\Big)$.

(c) *If $\Phi : \mathcal{X} \longrightarrow \mathcal{X}$ is equicontinuous and surjective, then $(\mathcal{X}, \Phi)$ is an odometer bundle.*

Proof. (a) For all $\mathsf{w} \in \mathbb{W}$, there is some R_{w} such that $\rho_{\mathsf{w}}(t) < R_{\mathsf{w}}$ for all $t \in \mathbb{N}$. Let $R := \max_{\mathsf{w} \in \mathbb{W}} R_{\mathsf{w}}$; then R is finite because $\mathbb{W}$ is finite. Let $\mathbb{U} := \mathbb{B}(\mathbb{W}, R)$.

Claim 1: *For all $\mathsf{w} \in \mathbb{W}$, and all $t \in \mathbb{N}$, we have $\Phi^t_{\text{in}}(\mathsf{w}) \subseteq \mathbb{U}$.*

Proof. (by contradiction) Suppose $\Phi^t_{\text{in}}(\mathsf{w}) \not\subseteq \mathbb{U}$. Let $\mathsf{v} \in \Phi^t_{\text{in}}(\mathsf{w}) \setminus \mathbb{U}$, and just as in the proof of Proposition 3.1 "$\Longleftarrow$", construct a path from v to w of length $L > R$. Conclude that $\rho_{\mathsf{w}}(t) > R$. Contradiction. ◇ `Claim 1`

Suppose $\mathbf{x}_{\mathbb{U}} = \mathbf{y}_{\mathbb{U}}$. Then for all $\mathsf{w} \in \mathbb{W}$, and all $t \in \mathbb{N}$, Claim 1 implies that $\Phi^t(\mathbf{x})_{\mathsf{w}} = \Phi^t(\mathbf{x})_{\mathsf{w}}$. In other words, $\Phi^t(\mathbf{y})_{\mathbb{W}} = \Phi^t(\mathbf{x})_{\mathbb{W}}$, for all $t \in \mathbb{N}$. Thus, $\mathbb{U}$ is an envelope for $\mathbb{W}$.

(b) "$\Longrightarrow$" follows immediately from part (a). For "$\Longleftarrow$", note that an equicontinuous system can have no sensitive points; now apply the contrapositive of Proposition 3.1(b).

(c) Let $\mathbb{W}_1 \subset \mathbb{W}_2 \subset \mathbb{W}_3 \subset \cdots$ be an increasing sequence of finite sets, with $\bigcup_{n=1}^{\infty} \mathbb{W}_n = \mathbb{V}$. For all $n \in \mathbb{N}$, let $\mathcal{B}_n := \mathcal{X}_{\mathbb{W}_n}$, and let $\Phi^{\mathbb{N}}_{\mathbb{W}_n} : \mathcal{X} \longrightarrow \mathcal{B}_n^{\mathbb{N}}$ be as in §2. Let $\mathcal{Y}_n := \Phi^{\mathbb{N}}_{\mathbb{W}_n}(\mathcal{X}) \subseteq \mathcal{B}_n^{\mathbb{N}}$. If $\sigma_n : \mathcal{B}_n^{\mathbb{N}} \longrightarrow \mathcal{B}_n^{\mathbb{N}}$ is the shift map, then $\sigma_n \circ \Phi^{\mathbb{N}}_{\mathbb{W}_n} = \Phi^{\mathbb{N}}_{\mathbb{W}_n} \circ \Phi$. Furthermore, $\sigma_n(\mathcal{Y}_n) = \mathcal{Y}_n$, because Φ is surjective.

For all $n \in \mathbb{N}$, let $\widetilde{\pi}_n : \mathcal{X}_{\mathbb{W}_{n+1}} \longrightarrow \mathcal{X}_{\mathbb{W}_n}$ be the projection (i.e. $\widetilde{\pi}_n(\mathbf{x}_{\mathbb{W}_{n+1}}) := \mathbf{x}_{\mathbb{W}_n}$ for all $\mathbf{x} \in \mathcal{X}$). Define $\pi_n : \mathcal{Y}_{n+1} \longrightarrow \mathcal{Y}_n$ as follows: if $\mathbf{y} \in \mathcal{Y}_{n+1}$, write $\mathbf{y} = [\mathbf{y}^t]_{t=0}^{\infty}$ where $\mathbf{y}^t \in \mathcal{X}_{\mathbb{W}_{n+1}}$ for all $t \in \mathbb{N}$; then define $\pi_n(\mathbf{y}) := [\widetilde{\pi}_n(\mathbf{y}^t)]_{t=0}^{\infty}$. Clearly, $\pi_n : (\mathcal{Y}_{n+1}, \sigma_{n+1}) \longrightarrow (\mathcal{Y}_n, \sigma_n)$ is a factor mapping, and $(\mathcal{X}, \Phi) = \lim \left\{ \cdots \xrightarrow{\pi_3} (\mathcal{Y}_3, \sigma_3) \xrightarrow{\pi_2} (\mathcal{Y}_2, \sigma_2) \xrightarrow{\pi_1} (\mathcal{Y}_1, \sigma_1) \right\}$.

Everything so far is true for any symbolic dynamical system. Now we use equicontinuity.

Claim 2: *For all $n \in \mathbb{N}$, $\mathcal{Y}_n$ is finite and $\sigma_n : \mathcal{Y}_n \longrightarrow \mathcal{Y}_n$ is a permutation.*

Proof. Let $\mathbb{U}_n \subset \mathbb{V}$ be the envelope of $\mathbb{W}_n$ (a finite set). For any $\mathbf{x}, \mathbf{x}' \in \mathcal{X}$, if $\mathbf{x}_{\mathbb{U}_n} = \mathbf{x}'_{\mathbb{U}_n}$, then $\Phi^{\mathbb{N}}_{\mathbb{W}_n}(\mathbf{x}) = \Phi^{\mathbb{N}}_{\mathbb{W}_n}(\mathbf{x}')$. Thus, $|\Phi^{\mathbb{N}}_{\mathbb{W}_n}(\mathcal{X})| \leq |\mathcal{X}_{\mathbb{U}_n}|$—in other words, $|\mathcal{Y}_n| \leq |\mathcal{X}_{\mathbb{U}_n}|$. But $|\mathcal{X}_{\mathbb{U}_n}|$ is finite because $\mathbb{U}_n$ is finite. Thus, $\mathcal{Y}_n$ is finite. Thus, σ_n is bijective (because it is surjective). ◇ `Claim 2`

Thus, we have represented $(\mathcal{X}, \Phi)$ as an inverse limit of finite permutation dynamical systems; thus, $(\mathcal{X}, \Phi)$ is an odometer bundle. □

For example: any symbolic dynamical system with the network in Figure 1 must be equicontinuous.

4 An expansive system of dimension two

The symmetry condition in Theorem 2.7 is probably not necessary. However, some sort of condition is required beyond merely superlinear connectivity. To demonstrate this, we will construct an example of a symbolic dynamical system which is posexpansive, despite having connectivity dimension two.



Figure 4: An expansive, two-dimensional symbolic dynamical system.

Let $\mathcal{A} := \mathbb{Z}_{/2} \times \mathbb{Z}_{/2}$, and let $\mathbb{V}$ be the digraph shown in Figure 4. Let $\mathbb{V}_\square$ be the set of vertices indicated by boxes and indexed by $\mathbb{M} := \{0, 2, 6, 12, 20, \ldots, m_k, \ldots\}$, where $m_k := \sum_{i=0}^{k} 2j$. We denote the m_kth square vertex by $\square_{m_k}$. Let $\mathbb{V}_\circ$ be the set of vertices indicated by circles; then $\mathbb{V} = \mathbb{V}_\square \sqcup \mathbb{V}_\circ$. We denote the nth element of $\mathbb{V}_\circ$ by $\bigcirc_n$.

For any vertex $\mathsf{v}_n \in \mathbb{V}$, the state of v_n is an ordered pair $\left(\begin{smallmatrix} a_n \\ b_n \end{smallmatrix}\right)$, where $a_n, b_n \in \mathbb{Z}_{/2}$. Let $\mathcal{X} := \left\{ \mathbf{a} \in \mathcal{A}^{\mathbb{V}} \; ; \; b_n = 0, \; \forall \; \mathsf{v}_n \in \mathbb{V}_\circ \right\}$. Thus, if $\mathcal{X}_\mathsf{v}$ is the projection of $\mathcal{X}$ onto vertex v, then we have $\mathcal{X}_\mathsf{v} = \mathbb{Z}_{/2} \times \mathbb{Z}_{/2}$ if $\mathsf{v} \in \mathbb{V}_\square$, and $\mathcal{X}_\mathsf{v} = \mathbb{Z}_{/2} \times \{0\}$ if $\mathsf{v} \in \mathbb{V}_\circ$. (Clearly $h(\mathcal{X}) \geq \log_2(2) = 1$, and $\mathcal{X}$ has weak independence.)

The local rule of each cell depends entirely upon its one or two input cells, and not on itself, as follows. For any $\bigcirc_n \in \mathbb{V}_\circ$, we define $\phi_n : \mathcal{X}_{n+1} \longrightarrow \mathcal{X}_n$ by $\phi_n \left(\begin{smallmatrix} a_{n+1} \\ b_{n+1} \end{smallmatrix}\right) = \left(\begin{smallmatrix} a_{n+1} \\ 0 \end{smallmatrix}\right)$ —that is, ϕ_n simply copies the first coordinate of $\bigcirc_{n+1}$ (or $\square_{n+1}$) into $\bigcirc_n$.

The cell $\square_{m_k}$ is connected to both $\bigcirc_{(m_k)+1}$ and $\square_{m_{(k+1)}}$. Its local rule $\phi_{m_k} : \mathcal{X}_{(m_k)+1} \times \mathcal{X}_{m_{(k+1)}} \longrightarrow \mathcal{X}_{m_k}$ is defined as follows:

$$\phi_{m_k} \left(\begin{pmatrix} a_{(m_k)+1} \\ 0 \end{pmatrix} , \begin{pmatrix} a_{m_{(k+1)}} \\ b_{m_{(k+1)}} \end{pmatrix} \right) \quad := \quad \begin{pmatrix} a_{(m_k)+1} \\ a_{m_{(k+1)}} + b_{m_{(k+1)}} \end{pmatrix} \tag{17}$$

Lemma 4.1 *The system $(\mathcal{X}, \Phi)$ is posexpansive*[1], *with posexpansive window $\{\square_0\}$.*

Proof. Define $\Gamma_1 : \mathcal{X} \longrightarrow {\mathbb{Z}_{/2}}^{\mathbb{N}}$ and $\Gamma_2 : \mathcal{X} \longrightarrow {\mathbb{Z}_{/2}}^{\mathbb{M}}$ by

$$\begin{aligned} \Gamma_1 \left[\begin{pmatrix} a_0 \\ b_0 \end{pmatrix}, \begin{pmatrix} a_1 \\ 0 \end{pmatrix}, \begin{pmatrix} a_2 \\ b_2 \end{pmatrix}, \begin{pmatrix} a_3 \\ 0 \end{pmatrix}, \begin{pmatrix} a_4 \\ 0 \end{pmatrix}, \begin{pmatrix} a_5 \\ 0 \end{pmatrix}, \begin{pmatrix} a_6 \\ b_6 \end{pmatrix}, \ldots \right] &:= [a_0, a_1, a_2, a_3, a_4, a_5, \ldots]; \\ \Gamma_2 \left[\begin{pmatrix} a_0 \\ b_0 \end{pmatrix}, \begin{pmatrix} a_1 \\ 0 \end{pmatrix}, \begin{pmatrix} a_2 \\ b_2 \end{pmatrix}, \begin{pmatrix} a_3 \\ 0 \end{pmatrix}, \begin{pmatrix} a_4 \\ 0 \end{pmatrix}, \begin{pmatrix} a_5 \\ 0 \end{pmatrix}, \begin{pmatrix} a_6 \\ b_6 \end{pmatrix}, \ldots \right] &:= [b_0, b_2, b_6, b_{12}, b_{20}, \ldots]. \end{aligned}$$

---

[1] See the start of §2 for the definitions of 'posexpansive' and 'window'.

Thus, $\Gamma := \Gamma_1 \times \Gamma_2 : \mathcal{X} \longrightarrow {\mathbb{Z}_{/2}}^{\mathbb{N}} \times {\mathbb{Z}_{/2}}^{\mathbb{M}}$ is a bijection. Let $\widetilde{\Phi} := \Gamma \circ \Phi \circ \Gamma^{-1} : {\mathbb{Z}_{/2}}^{\mathbb{N}} \times {\mathbb{Z}_{/2}}^{\mathbb{M}} \longrightarrow {\mathbb{Z}_{/2}}^{\mathbb{N}} \times {\mathbb{Z}_{/2}}^{\mathbb{M}}$; then Φ is conjugate to $\widetilde{\Phi}$ via Γ, so it suffices to show that the system $({\mathbb{Z}_{/2}}^{\mathbb{N}} \times {\mathbb{Z}_{/2}}^{\mathbb{M}}, \widetilde{\Phi})$ is posexpansive.

Let $(\mathbf{a}^0, \mathbf{b}^0) \in {\mathbb{Z}_{/2}}^{\mathbb{N}} \times {\mathbb{Z}_{/2}}^{\mathbb{M}}$ be some initial state, and let $(\mathbf{a}^t, \mathbf{b}^t) := \widetilde{\Phi}^t(\mathbf{a}^0, \mathbf{b}^0)$ for all $t \in \mathbb{N}$. Let $J \in \mathbb{N}$, and consider the sequence of the first m_J states of cell $\square_0$:

$$\mathbf{x}_J \quad := \quad \left(\begin{matrix} a_0^0 \\ b_0^0 \end{matrix}, \begin{matrix} a_0^1 \\ b_0^1 \end{matrix}, \ldots, \begin{matrix} a_0^{m_J} \\ b_0^{m_J} \end{matrix} \right).$$

Clearly, the information in $\mathbf{x}_J$ is sufficient to determine the values of $a_0^0, a_1^0, a_2^0, \ldots, a_{m_J}^0$, because for all $t \in \mathbb{N}$ we have $a_0^t = a_t^0$, because $\widetilde{\Phi}$ just acts like a unilateral shift on the 'a' coordinates. It remains to show that $\mathbf{x}_J$ is also sufficient to determine the values of $b_0^0, b_2^0, b_6^0, \ldots, b_{m_J}^0$.

Claim 1: *For any $j \in [1...J]$, and any $t \in [0 \ldots m_J - m_j]$, the information in $\mathbf{x}_J$ determines the value of $b_{m_j}^t$.*

Proof. (by induction on j)

Base case ($j = 1$). Note that $m_1 = 2$. We are given $b_0^0, b_0^1, \ldots, b_0^{m_J}$ in the bottom row of $\mathbf{x}_J$. For any $t \in [0 \ldots m_J - 2]$, we also know the value of $a_2^t = a_0^{t+2}$, and thus, we can compute $b_2^t = b_0^{t+1} - a_2^t$, because $b_0^{t+1} = a_2^t + b_2^t$, because substituting $m_0 = 0$ and $m_1 = 2$ into eqn.(17) yields

$$\begin{pmatrix} a_0^{t+1} \\ b_0^{t+1} \end{pmatrix} \quad = \quad \phi_0 \left(\begin{pmatrix} a_1^t \\ 0 \end{pmatrix}, \begin{pmatrix} a_2^t \\ b_2^t \end{pmatrix} \right) \quad := \quad \begin{pmatrix} a_1^t \\ a_2^t + b_2^t \end{pmatrix}.$$

Induction. Fix $j \in [1...J]$, and suppose we know the values of $b_{m_j}^t$ for all $t \in [0 \ldots m_J - m_j]$. Then we know $b_{m_j}^{t+1}$ for all $t \in \left[0 \ldots m_J - m_{(j+1)}\right]$ (because $m_J - m_{(j+1)} + 1 \leq m_J - m_j$ because $m_{(j+1)} > m_j + 1$). For any $t \in \left[0 \ldots m_J - m_{(j+1)}\right]$, we also know $a_{m_{(j+1)}}^t = a_0^{t+m_{(j+1)}}$ and thus, we can compute $b_{m_{(j+1)}}^t = b_{m_j}^{t+1} - a_{m_{(j+1)}}^t$, because $b_{m_j}^{t+1} = a_{m_{(j+1)}}^t + b_{m_{(j+1)}}^t$, because eqn.(17) says

$$\begin{pmatrix} a_{m_j}^{t+1} \\ b_{m_j}^{t+1} \end{pmatrix} \quad = \quad \phi_{m_j} \left(\begin{pmatrix} a_{m_j+1}^t \\ 0 \end{pmatrix}, \begin{pmatrix} a_{m_{(j+1)}}^t \\ b_{m_{(j+1)}}^t \end{pmatrix} \right) \quad := \quad \begin{pmatrix} a_{m_j+1}^t \\ a_{m_{(j+1)}}^t + b_{m_{(j+1)}}^t \end{pmatrix}.$$

$\diamondsuit$ `Claim 1`

In particular, Claim 1 implies that, for any $j \in [1...J]$, the information in $\mathbf{x}_J$ determines $b_{m_j}^0$. Thus, given $\mathbf{x}_J$, we can recover $a_0^0, a_2^0, a_3^0, \ldots, a_{m_J}^0$ and also $b_0^0, b_2^0, b_6^0, \ldots, b_{m_J}^0$. This works for all $J \in \mathbb{N}$; thus, $\{\square_0\}$ is a posexpansive window for $(\mathcal{X}, \Phi)$. □

Lemma 4.2 (a) *Φ has quadratic propagation*[2] $\left(\liminf_{t \to \infty} (\rho_{\mathsf{v}}(t)/t^2) > 0\right)$ *at every* $\mathsf{v} \in \mathbb{V}_{\square}$.

(b) $\dim_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\!\rightarrow) = 2$ *for all* $\mathsf{v} \in \mathbb{V}$.

---

[2] See equation (12) in §3 for the definition of 'propagation'.


$\Phi_{\rm in}^{\{0,1\}}(\square_0)$ $\Phi_{\rm in}^{[0\ldots 2]}(\square_0)$ $\Phi_{\rm in}^{[0\ldots 3]}(\square_0)$ $\Phi_{\rm in}^{[0\ldots 4]}(\square_0)$

$\Phi_{\rm in}^{[0\ldots 5]}(\square_0)$ $\Phi_{\rm in}^{[0\ldots 6]}(\square_0)$ $\Phi_{\rm in}^{[0\ldots 7]}(\square_0)$

Figure 5: The quadratic growth of $\Phi_{\rm in}^{[0\ldots T]}(\square_0)$.

Proof. (a) We will show this at vertex $\square_0$; the proof at other vertices is similar. Observe that $\Phi_{\rm in}^{[0\ldots T]}(\square_0)$ grows quadratically as $T \to \infty$, as shown in Figure 5. To be precise, for any $T > 0$, $\Phi_{\rm in}^{[0\ldots T]}(\square_0)$ contains the cells $\{\square_0, \square_2, \square_6, \ldots, \square_{m_T}\}$, and also contains the cells $\bigcirc_{m_t+s}$ for each $t \in [1\ldots T]$ and $s \in [1\ldots T-t]$.

(b) From (a) we have $\dim_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\rightarrow) \geq 2$ for all $\mathsf{v} \in \mathbb{V}_\square$ (because $|\mathbb{B}(\mathsf{v}, r)| \geq \rho_{\mathsf{v}}(r)$ for all $r \in \mathbb{N}$). But every vertex is downstream from some element of $\mathbb{V}_\square$; thus, Lemma 1.7 implies that $\dim_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\rightarrow) \geq 2$ for all $\mathsf{v} \in \mathbb{V}$. On the other hand, it is easy to see that $\dim_{\mathsf{v}}(\mathbb{V}, \bullet\!\!\rightarrow) \leq 2$. $\square$

5 Lipschitz metrics on Cantor systems

The counterexample of §4 shows that connectivity dimension is *not* invariant under topological conjugacy: some systems with dimension two are conjugate to subshifts of $(\mathcal{A}^{\mathbb{N}}, \sigma)$, and the system $(\mathcal{A}^{\mathbb{N}}, \sigma)$ has dimension one. (Likewise, §4 shows that the growth rate of the propagation function ρ is not a conjugacy invariant.) However, we will now show that connectivity dimension *is* invariant under a slightly refined notion of conjugacy, once we impose a suitable metric structure on the pattern space $\mathcal{X}$ (see Corollary 5.8 below).

Let $\mathcal{X}$ be any Cantor space, and let $d : \mathcal{X}^2 \longrightarrow \mathbb{R}_+$ be a metric compatible with the Cantor topology on $\mathcal{X}$. The pair $(\mathcal{X}, d)$ will be called a *Cantor metric space*. Let $\Phi : \mathcal{X} \longrightarrow \mathcal{X}$

be any continuous self-map. We say Φ is *d-Lipschitz* if there is a constant $\lambda > 0$ such that, for any $\mathbf{x}, \mathbf{x}' \in \mathcal{X}$,

$$d\left(\Phi(\mathbf{x}), \Phi(\mathbf{x}')\right) \quad \leq \quad \lambda \cdot d(\mathbf{x}, \mathbf{x}'). \tag{18}$$

In this case, d is called a ***Lipschitz metric*** for Φ. The smallest λ satisfying eqn.(18) is called the *d-**Lipschitz constant*** of Φ. More generally, a ***Lipschitz pseudometric*** is a pseudometric $d : \mathcal{X}^2 \longrightarrow \mathbb{R}_+$ satisfying eqn.(18).

Example 5.1. Let $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ be a symbolic dynamical system, with network $(\mathbb{V}, \bullet\!\!\!\rightarrow)$. For any $\mathsf{v} \in \mathbb{V}$ and $\lambda > 1$, we define the pseudometric $d_{\mathsf{v},\lambda} : \mathcal{A}^{\mathbb{V}} \times \mathcal{A}^{\mathbb{V}} \longrightarrow \mathbb{R}_+$ as follows: for all $\mathbf{a}, \mathbf{b} \in \mathcal{A}^{\mathbb{V}}$,

$$d_{\mathsf{v},\lambda}(\mathbf{a}, \mathbf{b}) \quad := \quad \frac{1}{\lambda^{R(\mathbf{a},\mathbf{b})}}, \quad \text{where} \quad R(\mathbf{a}, \mathbf{b}) \quad := \quad \max\left\{r \in \mathbb{N} \; ; \; \mathbf{a}_{\mathbb{B}(\mathsf{v},r)} = \mathbf{b}_{\mathbb{B}(\mathsf{v},r)}\right\}. \tag{19}$$

Then $d_{\mathsf{v},\lambda}$ is a Φ-Lipschitz pseudometric with constant λ. To see this, let $\mathbf{x}, \mathbf{x}' \in \mathcal{X}$. If $R(\mathbf{x}, \mathbf{x}') = r$, then $d(\mathbf{x}, \mathbf{x}') = \frac{1}{\lambda^r}$. But if $R(\mathbf{x}, \mathbf{x}') = r$, then $R\left[\Phi(\mathbf{x}'), \Phi(\mathbf{x}')\right] \geq r - 1$, so $d\left[\Phi(\mathbf{x}'), \Phi(\mathbf{x}')\right] \leq \frac{1}{\lambda^{r-1}} = \lambda \cdot d(\mathbf{x}, \mathbf{x}')$, as desired. ♢

The pseudometric $d_{\mathsf{v},\lambda}$ in Example 5.1 is not necessarily a true metric, unless $\bigcup_{r=0}^{\infty} \mathbb{B}(\mathsf{v}, r) = \mathbb{V}$, which is not the case unless v is downstream from every element of v. For many digraphs, there is no vertex with this property. Instead, let us say that a subset $\mathbb{U} \subset \mathbb{V}$ is an ***estuary*** if, for every $\mathsf{v} \in \mathbb{V}$, there exists some $\mathsf{u} \in \mathbb{U}$ with $\mathsf{v} \rightsquigarrow \mathsf{u}$. For example, $\mathbb{V}$ itself is an estuary. If $(\mathbb{V}, \bullet\!\!\!\rightarrow)$ is biconnected, then any nonempty subset of $\mathbb{V}$ (even a singleton) is an estuary. More generally, if $\mathbb{U}$ contains at least one vertex from each biconnected component of $\mathbb{V}$, then $\mathbb{U}$ is an estuary.

Example 5.2. Let $\mathbb{U} \subset \mathbb{V}$ be an estuary. Let $\mathbf{c} := (c_{\mathsf{u}})_{\mathsf{u}\in\mathbb{U}} \in \mathbb{R}_+^{\mathbb{U}}$ be a sequence of positive ***coefficients*** for the elements in $\mathbb{U}$, such that $\sum\limits_{\mathsf{u}\in\mathbb{U}} c_{\mathsf{u}} < \infty$ (this is possible because $\mathbb{U}$ is always countable, because $\mathbb{V}$ is countable). Fix $\lambda > 1$, and for all $\mathsf{u} \in \mathbb{U}$, let $d_{\mathsf{u},\lambda}$ be the pseudometric from Example 5.1. Define the metric $d_{\mathbf{c},\lambda} : \mathcal{X} \times \mathcal{X} \longrightarrow \mathbb{R}_+$ by

$$d_{\mathbf{c},\lambda}(\mathbf{a}, \mathbf{b}) \quad := \quad \sum_{\mathsf{u}\in\mathbb{U}}^{\infty} c_{\mathsf{u}}\, d_{\mathsf{u},\lambda}(\mathbf{a}, \mathbf{b}).$$

Then $d_{\mathbf{c},\lambda}$ is a true metric, because $\bigcup_{r=0}^{\infty} \mathbb{B}(\mathbb{U}, r) = \mathbb{V}$, because $\mathbb{U}$ is an estuary. Also, $d_{\mathbf{c},\lambda}$ satisfies eqn.(18), because each of the pseudometrics $d_{\mathsf{u},\lambda}$ satisfies eqn.(18) (by Example 5.1). Thus, $d_{\mathbf{c},\lambda}$ is a Lipschitz metric for Φ. We say that $d_{\mathbf{c},\lambda}$ is ***based at*** $\mathbb{U}$. ♢

Observe that the metric on $\mathcal{A}^{\mathbb{V}}$ in Example 5.2 can be defined for any digraph structure on $(\mathbb{V}, \bullet\!\!\!\rightarrow)$ (without reference to any particular map $\Phi : \mathcal{A}^{\mathbb{V}} \longrightarrow \mathcal{A}^{\mathbb{V}}$). Any Cantor dynamical system can be represented as a symbolic dynamical system, so Example 5.2 shows that any Cantor dynamical system admits a Lipschitz metric. Indeed, it admits many such metrics, because the Lipschitz constant λ, the estuary $\mathbb{U}$, and the coefficient system $\mathbf{c}$ in Example 5.2 can be chosen arbitrarily.

Dimension and Entropy. Let $(\mathcal{X}, d)$ be a metric space. For any $\epsilon > 0$, a *open ϵ-cover* is a covering of $\mathcal{X}$ by open sets whose diameters are each at most ϵ. Let $N_\epsilon(\mathcal{X})$ be the minimal cardinality of any open ϵ-cover of $(\mathcal{X}, d)$. We define

$$\begin{aligned} \underline{\dim}(\mathcal{X}, d) &:= \liminf_{\epsilon \to 0} \frac{\log_\alpha \left[\log_\beta(N_\epsilon(\mathcal{X}))\right]}{\log_\alpha \left[-\log_\gamma(\epsilon)\right]} \\ \text{and} \qquad \overline{\dim}(\mathcal{X}, d) &:= \limsup_{\epsilon \to 0} \frac{\log_\alpha \left[\log_\beta(N_\epsilon(\mathcal{X}))\right]}{\log_\alpha \left[-\log_\gamma(\epsilon)\right]}. \end{aligned} \tag{20}$$

(Here, $\alpha, \beta, \gamma > 1$ are any constants, and need not be equal —the limits in (20) are independent of the choice of α, β, γ). If $\underline{\dim}(\mathcal{X}, d) = \overline{\dim}(\mathcal{X}, d)$, then we refer to their common value as "$\dim(\mathcal{X}, d)$", the *dimension* of $(\mathcal{X}, d)$. Note that formula (20) differs from the 'box-counting dimension' $\operatorname{boxdim}(\mathcal{X}, d) := \lim_{\epsilon \to 0} \frac{\log_2(N_\epsilon)}{-\log_2(\epsilon)}$ by the extra logarithms. Also, $\dim(\mathcal{X}, d)$ is meaningful even when $\operatorname{boxdim}(\mathcal{X}, d) = \infty$. (Indeed, $\operatorname{boxdim}(\mathcal{X}, d)$ is finite iff $\dim(\mathcal{X}, d) = 1$). We will now show that $\dim(\mathcal{X}, d)$ is closely related to the 'connectivity dimension' of the digraph $(\mathbb{V}, \bullet\!\!\to)$.

Let $\underline{\dim}(\mathbb{V}, \bullet\!\!\to)$ and $\overline{\dim}(\mathbb{V}, \bullet\!\!\to)$ be as defined in eqn.(1) of §1. If $\mathbb{U} \subseteq \mathbb{V}$ is an estuary for $\mathbb{V}$, then Lemma 1.7 implies that

$$\overline{\dim}(\mathbb{V}, \bullet\!\!\to) \quad = \quad \sup_{\mathsf{u} \in \mathbb{U}} \limsup_{r \to \infty} \frac{\log |\mathbb{B}(\mathsf{u}, r)|}{\log(r)}. \tag{21}$$

We say $(\mathbb{V}, \bullet\!\!\to)$ has *uniform dimension* on $\mathbb{U}$ if the 'sup' and 'limsup' can be exchanged:

$$\overline{\dim}(\mathbb{V}, \bullet\!\!\to) \quad = \quad \limsup_{r \to \infty} \sup_{\mathsf{u} \in \mathbb{U}} \frac{\log |\mathbb{B}(\mathsf{u}, r)|}{\log(r)}. \tag{22}$$

Heuristically, eqn.(22) means that the limsups in eqn.(21) converge 'uniformly' on $\mathbb{U}$. For example, if $\mathbb{U}$ is finite, then $(\mathbb{V}, \bullet\!\!\to)$ automatically has uniform dimension on $\mathbb{U}$.

A nonnegative sequence $\{c_j\}_{j=1}^\infty$ has *precipitous decay* if

$$\lim_{\epsilon \to 0} \frac{\ln[J(\epsilon)]}{\ln|\ln(\epsilon)|} \quad = \quad 0, \quad \text{where, for all } \epsilon > 0, \quad J(\epsilon) \quad := \quad \min\left\{ J \in \mathbb{N} \, ; \, \sum_{j=J+1}^\infty c_j < \frac{\epsilon}{2} \right\}. \tag{23}$$

Let $\mathbb{U} \subseteq \mathbb{V}$ and let $\mathbf{c} = (c_\mathsf{u})_{\mathsf{u} \in \mathbb{U}}$ be some coefficient sequence. Suppose we enumerate $\mathbb{U}$ as $\mathbb{U} = \{\mathsf{u}_j\}_{j=0}^\infty$; then we can define $c'_j := c_{\mathsf{u}_j}$ for all $j \in \mathbb{N}$; then we say $\mathbf{c}$ has *precipitous decay* if the sequence $\{c'_j\}_{j=1}^\infty$ has precipitous decay.

Example 5.3. (a) If $\{c_j\}_{j=1}^\infty$ has only finitely many nonzero terms, then $\{c_j\}_{j=1}^\infty$ has precipitous decay. (*Proof.* If $c_j = 0$ for all $j \geq J_0$, then $J(\epsilon) \leq J_0$ for all ϵ.)

(b) Let $c_j := \exp(-e^j) \cdot e^j$ for all $j \in \mathbb{N}$. Then $\{c_j\}_{j=1}^\infty$ has precipitous decay. (*Proof.* If $F(x) := -\exp(-e^x)$, then $F'(x) = \exp(-e^x) \cdot e^x$, so $c_j = F'(j)$ for all $j \in \mathbb{N}$. Thus, $\sum_{j=J+1}^\infty c_j < \int_{J+1}^\infty F'(x)\, dx = -F(J+1)$, so $J(\epsilon) \leq F^{-1}(-\epsilon/2) = \ln[-\ln(\epsilon/2)]$.) ♢

Proposition 5.4 *Let* $(\mathbb{V}, \bullet\!\!\to)$ *be a digraph, let* $\mathbb{U} \subset \mathbb{V}$ *be an estuary, let* $\lambda > 0$*, and let* $d = d_{\lambda,\mathbf{c}} : \mathcal{X}^2 \longrightarrow \mathbb{R}_+$ *be a metric based on* $\mathbb{U}$*, as in* Example 5.2. *Let* $\mathcal{X} \subseteq \mathcal{A}^{\mathbb{V}}$ *be a pattern space.*

(a) *Suppose* $\mathbb{U}' := \{\mathsf{u} \in \mathbb{U} \ ; \ \underline{h}_{\mathsf{u}}(\mathcal{X}) > 0\}$ *is nonempty, and let* $\underline{D} := \sup_{\mathsf{u}\in\mathbb{U}'} \underline{\dim}_{\mathsf{u}}(\mathbb{V}, \bullet\!\!\to)$*. Then* $\underline{\dim}(\mathcal{X}, d) \geq \underline{D}$*. In particular,* $\underline{\dim}(\mathcal{X}, d) \geq \underline{\dim}(\mathbb{V}, \bullet\!\!\to)$*.*

Let $\mathbf{c} = (c_{\mathsf{u}})_{\mathsf{u}\in\mathbb{U}}$ *be the coefficients used to define* d*.*

(b) *If* $\mathbf{c}$ *has precipitous decay and* $(\mathbb{V}, \bullet\!\!\longrightarrow)$ *has uniform dimension on* $\mathbb{U}$*, then* $\overline{\dim}(\mathcal{X}, d) \ \leq \ \overline{\dim}(\mathbb{V}, \bullet\!\!\to)$*.*

(c) *In particular, if* $\mathbb{U}$ *is finite, then* $\overline{\dim}(\mathcal{X}, d) \ \leq \ \overline{\dim}(\mathbb{V}, \bullet\!\!\to)$*.*

Proof. Let $\mathbb{W} \subset \mathbb{V}$ be any finite set. For all $\mathbf{w} \in \mathcal{X}_{\mathbb{W}}$, let $\langle\mathbf{w}\rangle := \{\mathbf{x} \in \mathcal{X} \ ; \ \mathbf{x}_{\mathbb{W}} = \mathbf{w}\}$ be the *cylinder set* defined by $\mathbf{w}$. The collection $\mathcal{C}_{\mathbb{W}} := \{\langle\mathbf{w}\rangle \ ; \ \mathbf{w} \in \mathcal{X}_{\mathbb{W}}\}$ is an open cover of $\mathcal{X}$.

(a) Let $\underline{\delta} < \underline{D}$.

Claim 1: *There exists* $\epsilon_1 > 0$*,* $H > 0$*, and* $L \in \mathbb{R}$ *such that, for all* $\epsilon \in (0, \epsilon_1)$*, we have*

$$\ln\left(\log_2[N_\epsilon(\mathcal{X})]\right) \quad > \quad \ln(H) + \underline{\delta} \cdot \ln(L - \log_\lambda(\epsilon)).$$

Proof. For any $\epsilon > 0$, let $\mathbb{U}(\epsilon) := \{\mathsf{u} \in \mathbb{U} \ ; \ c_{\mathsf{u}} > \epsilon\}$ (which is finite because $\mathbf{c}$ is summable). For all $\mathsf{u} \in \mathbb{U}(\epsilon)$, let $r_{\mathsf{u}}(\epsilon) := \lfloor \log_\lambda(c_{\mathsf{u}}/\epsilon) \rfloor$. Let $\mathbb{W}(\epsilon) := \bigcup_{\mathsf{u}\in\mathbb{U}(\epsilon)} \mathbb{B}(\mathsf{u}, r_{\mathsf{u}}(\epsilon))$.

Claim 1.1: *Let* $\mathbf{x}, \mathbf{y} \in \mathcal{X}$*. If* $\mathbf{x}_{\mathbb{W}(\epsilon)} \neq \mathbf{y}_{\mathbb{W}(\epsilon)}$*, then* $d(\mathbf{x}, \mathbf{y}) > \epsilon$*.*

Proof. If $\mathbf{x}_{\mathbb{W}(\epsilon)} \neq \mathbf{y}_{\mathbb{W}(\epsilon)}$, then there exists $\mathsf{u} \in \mathbb{U}(\epsilon)$ with $\mathbf{x}_{\mathbb{B}(\mathsf{u},r_{\mathsf{u}}(\epsilon))} \neq \mathbf{y}_{\mathbb{B}(\mathsf{u},r_{\mathsf{u}}(\epsilon))}$. Thus

$$d(\mathbf{x}, \mathbf{y}) \quad := \quad \sum_{\mathsf{v}\in\mathbb{U}} c_{\mathsf{v}}\, d_{\mathsf{v},\lambda}(\mathbf{x}, \mathbf{y}) \quad \geq \quad c_{\mathsf{u}} \cdot d_{\mathsf{u},\lambda}(\mathbf{x}, \mathbf{y}) \quad \underset{(*)}{>} \quad \frac{c_{\mathsf{u}}}{\lambda^{r_u(\epsilon)}} \quad \underset{(\dagger)}{\geq} \quad \frac{c_{\mathsf{u}}\,\epsilon}{c_{\mathsf{u}}} \quad = \quad \epsilon,$$

as desired. Here, $(*)$ is by eqn.(19), and $(\dagger)$ is because $r_u(\epsilon) \leq \log_\lambda(c_{\mathsf{u}}/\epsilon)$. ∇ `Claim 1.1`

By hypothesis, there exists $\mathsf{u}^* \in \mathbb{U}'$ with $\underline{\dim}_{\mathsf{u}^*}(\mathbb{B}, \bullet\!\!\to) > \underline{\delta}$. Now, $\lim_{\epsilon\to 0} \mathbb{U}(\epsilon) = \mathbb{U}$ (because $c_{\mathsf{u}} > 0$ for all $\mathsf{u} \in \mathbb{U}$), so there exists $\epsilon_0 > 0$ such that, if $\epsilon \in (0, \epsilon_0)$, then $\mathsf{u}^* \in \mathbb{U}(\epsilon)$. Let $0 < H < \underline{h}_{\mathsf{u}^*}(\mathcal{X})$. Defining equations (1) and (2) in §1 say there exists $R > 0$ such that, for all $r > R$, we have $\dfrac{\ln|\mathbb{B}(\mathsf{u}^*, r)|}{\ln(r)} > \underline{\delta}$ and $\dfrac{\log_2\left|\mathcal{X}_{\mathbb{B}(\mathsf{u}^*,r)}\right|}{|\mathbb{B}(\mathsf{u}^*, r)|} > H$. But $\lim_{\epsilon\to 0} r_{\mathsf{u}^*}(\epsilon) = \infty$. Thus, there exists $\epsilon_1 \in (0, \epsilon_0)$ such that, if $\epsilon \in (0, \epsilon_1)$, then $r_{\mathsf{u}^*}(\epsilon) > R$; hence

$$\ln|\mathbb{B}(\mathsf{u}, r_{\mathsf{u}^*}(\epsilon))| \quad > \quad \underline{\delta} \cdot \ln(r_{\mathsf{u}^*}(\epsilon)) \tag{24}$$

$$\text{and} \quad \log_2\left|\mathcal{X}_{\mathbb{B}(\mathsf{u}^*,r_{\mathsf{u}^*}(\epsilon))}\right| \quad > \quad H \cdot |\mathbb{B}(\mathsf{u}^*, r_{\mathsf{u}^*}(\epsilon))|\,. \tag{25}$$

Let $\epsilon \in (0, \epsilon_1)$, and let $\mathcal{C}_\epsilon$ be a minimal open ϵ-cover; then Claim 1.1 implies that each cell of $\mathcal{C}_\epsilon$ can intersect at most one cylinder set from the cover $\mathcal{C}_{\mathbb{W}(\epsilon)}$. Thus,

$$\begin{aligned}
N_\epsilon(\mathcal{X}) \quad &= \quad |\mathcal{C}_\epsilon| \quad \geq \quad |\mathcal{C}_{\mathbb{W}(\epsilon)}| \quad = \quad |\mathcal{X}_{\mathbb{W}(\epsilon)}| \quad \geq \quad \max_{\mathsf{u} \in \mathbb{U}(\epsilon)} \left|\mathcal{X}_{\mathbb{B}(\mathsf{u}, r_{\mathsf{u}}(\epsilon))}\right| \\
&\underset{(*)}{\geq} \quad \left|\mathcal{X}_{\mathbb{B}(\mathsf{u}^*, r_{\mathsf{u}^*}(\epsilon))}\right|. \\
\text{Thus,} \quad \log_2[N_\epsilon(\mathcal{X})] \quad &\geq \quad \log_2 \left|\mathcal{X}_{\mathbb{B}(\mathsf{u}^*, r_{\mathsf{u}^*}(\epsilon))}\right| \quad \underset{(\dagger)}{>} \quad H \cdot |\mathbb{B}(\mathsf{u}^*, r_{\mathsf{u}^*}(\epsilon))| \\
\text{Thus,} \quad \ln\left(\log_2[N_\epsilon(\mathcal{X})]\right) \quad &> \quad \ln(H) + \ln|\mathbb{B}(\mathsf{u}^*, r_{\mathsf{u}^*}(\epsilon))| \\
&\underset{(\ddagger)}{>} \quad \ln(H) + \underline{\delta} \cdot \ln(r_{\mathsf{u}^*}(\epsilon)) \quad \underset{(\diamond)}{>} \quad \ln(H) + \underline{\delta} \cdot \ln(\log_\lambda(c_{\mathsf{u}^*}/\epsilon) - 1) \\
&= \quad \ln(H) + \underline{\delta} \cdot \ln(L - \log_\lambda(\epsilon)),
\end{aligned}$$

where $L := \log(c_{\mathsf{u}^*}) - 1$. Here, $(*)$ is because $\mathsf{u}^* \in \mathbb{U}(\epsilon)$ because $\epsilon < \epsilon_0$. $(\dagger)$ is by (25), $(\ddagger)$ is by (24), and $(\diamond)$ is because $r_{\mathsf{u}^*}(\epsilon) := \lfloor \log_\lambda(c_{\mathsf{u}^*}/\epsilon) \rfloor > \log_\lambda(c_{\mathsf{u}^*}/\epsilon) - 1$. ◇ `Claim 1`

$$\begin{aligned}
\text{We now have} \qquad \underline{\dim}(\mathcal{X}, d) \quad &\underset{(*)}{:=} \quad \liminf_{\epsilon \to 0} \frac{\ln\left[\log_2(N_\epsilon(\mathcal{X}))\right]}{\ln\left[-\log_\lambda(\epsilon)\right]} \\
&\underset{(\dagger)}{\geq} \quad \liminf_{\epsilon \to 0} \frac{\ln(H) + \underline{\delta} \cdot \ln(L - \log_\lambda(\epsilon))}{\ln\left[-\log_\lambda(\epsilon)\right]} \quad = \quad \underline{\delta},
\end{aligned}$$

where $(*)$ is by setting $\alpha := e$, $\beta := 2$, and $\gamma := \lambda$ in definition (20), while $(\dagger)$ is by Claim 1. This holds for any $\underline{\delta} < \underline{D}$. Thus, we conclude that $\underline{\dim}(\mathcal{X}, d) \geq \underline{D}$, as desired.

(b) Fix some enumeration $\mathbb{U} = \{\mathsf{u}_j\}_{j=0}^\infty$ and define $c_j := c_{\mathsf{u}_j}$ for all $j \in \mathbb{N}$. For all $\epsilon > 0$, let $J(\epsilon)$ be as in eqn.(23). Let $\overline{\delta} > \overline{\dim}(\mathbb{V}, \bullet\!\!\rightarrow)$. Let $S := \sum_{j=0}^\infty c_j < \infty$.

Claim 2: *There exists* $\epsilon_1 > 0$ *and constants* $L_1, L_2 > 0$ *such that, for any* $\epsilon \in (0, \epsilon_1)$*:*

$$\ln\left(\log_2[N_\epsilon(\mathcal{X})]\right) \quad \leq \quad \ln[J(\epsilon)] \; + \; L_1 \; + \; \overline{\delta} \cdot \ln[L_2 - \log_\lambda(\epsilon)].$$

Proof. For any $\epsilon > 0$, let $r(\epsilon) := \lceil \log_\lambda(2S/\epsilon) \rceil$, and let $\mathbb{W}(\epsilon) := \bigcup_{j=0}^{J(\epsilon)} \mathbb{B}[\mathsf{v}_j, r(\epsilon)]$.

Claim 2.1: *Let* $\mathbf{x}, \mathbf{y} \in \mathcal{X}$*. If* $\mathbf{x}_{\mathbb{W}(\epsilon)} = \mathbf{y}_{\mathbb{W}(\epsilon)}$*, then* $d(\mathbf{x}, \mathbf{y}) < \epsilon$.

Proof. We have

$$\begin{aligned}
d(\mathbf{x}, \mathbf{y}) \quad &:= \quad \sum_{j=0}^\infty c_j \, d_{\mathsf{v}_j, \lambda}(\mathbf{x}, \mathbf{y}) \quad = \quad \sum_{j=0}^{J(\epsilon)} c_j \, d_{\mathsf{v}_j, \lambda}(\mathbf{x}, \mathbf{y}) \; + \sum_{j=J(\epsilon)+1}^\infty c_j \, d_{\mathsf{v}_j, \lambda}(\mathbf{x}, \mathbf{y}) \\
&\leq \quad \left(\sum_{j=0}^{J(\epsilon)} c_j\right) \cdot \max_{0 \leq j \leq J(\epsilon)} \left(d_{\mathsf{v}_j, \lambda}(\mathbf{x}, \mathbf{y})\right) \; + \; \left(\sum_{j=J(\epsilon)+1}^\infty c_j\right) \cdot \max_{j \geq J(\epsilon)} \left(d_{\mathsf{v}_j, \lambda}(\mathbf{x}, \mathbf{y})\right) \\
&\underset{(*)}{\leq} \quad \frac{S}{\lambda^{r(\epsilon)}} \; + \; \frac{\epsilon}{2} \cdot 1 \quad \underset{(\dagger)}{\leq} \quad \frac{\epsilon S}{2S} \; + \; \frac{\epsilon}{2} \quad = \quad \frac{\epsilon}{2} + \frac{\epsilon}{2} \quad = \quad \epsilon, \qquad \text{as desired.}
\end{aligned}$$

Here, $(*)$ is because $d_{\mathsf{v}_j,\lambda}(\mathbf{x},\mathbf{y}) \leq \frac{1}{\lambda^{r(\epsilon)}}$ for all $j \in [0...J(\epsilon)]$ because $\mathbf{x}_{\mathbb{B}[\mathsf{v}_j,r(\epsilon)]} = \mathbf{x}_{\mathbb{B}[\mathsf{v}_j,r(\epsilon)]}$ for all $j \in [0...J(\epsilon)]$; meanwhile, $\displaystyle\sum_{j=0}^{J(\epsilon)} c_j \leq \sum_{j=0}^{\infty} c_j = S$, and $\displaystyle\sum_{j=J(\epsilon)+1}^{\infty} c_j < \frac{\epsilon}{2}$ by definition of $J(\epsilon)$. Finally, $(\dagger)$ is because $r = \lceil \log_\lambda(2S/\epsilon) \rceil$. $\triangledown$ `Claim 2.1`

Now, $\overline{\delta} > \overline{\dim}(\mathbb{V}, \bullet\!\!\rightarrow)$, so eqn.(22) yields some $R \in \mathbb{N}$ such that, for all $r \in \mathbb{N}$:

$$\text{if } r > R, \text{ then} \qquad \sup_{\mathsf{u}\in\mathbb{U}} \ln|\mathbb{B}(\mathsf{u},r)| \quad < \quad \overline{\delta}\cdot\ln(r). \tag{26}$$

Now, $\lim\limits_{\epsilon\to 0} r(\epsilon) = \infty$, so there exists $\epsilon_1 > 0$ such that, if $\epsilon \in (0,\epsilon_1)$, then $r(\epsilon) > R$. Let $\epsilon \in (0,\epsilon_1)$. Claim 2.1 implies that $\mathcal{C}_{\mathbb{W}(\epsilon)}$ is an ϵ-open cover of $\mathcal{X}$. Thus,

$$N_\epsilon(\mathcal{X}) \;\leq\; |\mathcal{C}_{\mathbb{W}(\epsilon)}| \;=\; |\mathcal{X}_{\mathbb{W}(\epsilon)}| \;\leq\; \prod_{j=0}^{J(\epsilon)} \left|\mathcal{X}_{\mathbb{B}(\mathsf{v}_j,r(\epsilon))}\right|.$$

$$\begin{aligned}
\text{Thus,}\quad \log_2[N_\epsilon(\mathcal{X})] &\leq \sum_{j=0}^{J(\epsilon)} \log_2\left|\mathcal{X}_{\mathbb{B}(\mathsf{v}_j,r(\epsilon))}\right| \;\leq\; \sum_{j=0}^{J(\epsilon)} (\log_2|\mathcal{A}|)\cdot\ |\mathbb{B}(\mathsf{v}_j,r(\epsilon))| \\
&\leq J(\epsilon)\cdot\log_2|\mathcal{A}|\cdot \max_{0\leq j\leq J(\epsilon)} |\mathbb{B}(\mathsf{v}_j,r(\epsilon))| \\
&\leq J(\epsilon)\cdot\log_2|\mathcal{A}|\cdot \sup_{\mathsf{u}\in\mathbb{U}} |\mathbb{B}(\mathsf{u},r(\epsilon))|. \\
\text{Thus,}\quad \ln(\log_2[N_\epsilon(\mathcal{X})]) &\leq \ln[J(\epsilon)] \;+\; \ln(\log_2|\mathcal{A}|) \;+\; \sup_{\mathsf{u}\in\mathbb{U}} \ln|\mathbb{B}(\mathsf{u},r(\epsilon))| \\
&\underset{(*)}{\leq} \ln[J(\epsilon)] \;+\; \ln(\log_2|\mathcal{A}|) \;+\; \overline{\delta}\cdot\ln(r(\epsilon)) \\
&\underset{(\dagger)}{\leq} \ln[J(\epsilon)] \;+\; L_1 \;+\; \overline{\delta}\cdot\ln(r(\epsilon)) \\
&\underset{(\ddagger)}{\leq} \ln[J(\epsilon)] \;+\; L_1 \;+\; \overline{\delta}\cdot\ln[1+\log_\lambda(2S/\epsilon)] \\
&\leq \ln[J(\epsilon)] \;+\; L_1 \;+\; \overline{\delta}\cdot\ln[L_2 - \log_\lambda(\epsilon)],
\end{aligned}$$

where $L_2 := 1+\log_\lambda(2S)$. Here, $(*)$ is by eqn.(26), because $r(\epsilon) > R$ because $\epsilon \in (0,\epsilon_1)$. In $(\dagger)$ we define $L_1 := \ln(\log_2|\mathcal{A}|)]$. Finally, $(\ddagger)$ is because $r := \lceil \log_\lambda(2S/\epsilon) \rceil \leq 1 + \log_\lambda(2S/\epsilon)$. $\diamondsuit$ `Claim 2`

$$\begin{aligned}
\text{We now have}\qquad \overline{\dim}(\mathcal{X},d) &\underset{(*)}{:=} \limsup_{\epsilon\to 0} \frac{\ln[\log_2(N_\epsilon(\mathcal{X}))]}{\ln[-\log_\lambda(\epsilon)]} \\
&\underset{(\dagger)}{\leq} \limsup_{\epsilon\to 0} \frac{\ln[J(\epsilon)] \;+\; L_1 \;+\; \overline{\delta}\cdot\ln[L_2 - \log_\lambda(\epsilon)]}{\ln[-\log_\lambda(\epsilon)]} \\
&= \overline{\delta} \;+\; \lim_{\epsilon\to 0} \frac{\ln[J(\epsilon)]}{\ln[-\log_\lambda(\epsilon)]} \;\underset{(\ddagger)}{=}\; \overline{\delta}.
\end{aligned}$$

Here, $(*)$ is by setting $\alpha := e$, $\beta := 2$, and $\gamma := \lambda$ in definition (20), and $(\dagger)$ is by Claim 2. Meanwhile, $(\ddagger)$ is because $\{c_j\}_{j=1}^\infty$ has precipitous decay.

Thus works for any $\overline{\delta} > \overline{\dim}(\mathbb{X}, \bullet\!\!\to)$; we conclude that $\overline{\dim}(\mathcal{X}, d) \le \overline{\dim}(\mathbb{X}, \bullet\!\!\to)$.

(c) follows immediately from (b), because if $\mathbb{U}$ is finite, then clearly $(\mathbb{V}, \bullet\!\!\to)$ has uniform dimension on $\mathbb{U}$, and $\mathbf{c}$ has precipitous decay. ☐

Let $(\mathbb{V}, \bullet\!\!\to)$ be a dimensionally homogeneous digraph [i.e. $\underline{\dim}(\mathbb{V}, \bullet\!\!\to) = \overline{\dim}(\mathbb{V}, \bullet\!\!\to)$]. If $\mathcal{X} \subseteq \mathcal{A}^{\mathbb{V}}$ is a pattern space, and $d : \mathcal{X}^2 \longrightarrow \mathbb{R}_+$ is a Cantor metric, then we say that d is *dimensionally compatible* if $\dim(\mathcal{X}, d) = \dim(\mathbb{V}, \bullet\!\!\to)$. Proposition 5.4 suggests that for 'most' dimensionally homogeneous digraphs, any pattern space with nonzero entropy admits a dimensionally compatible metric. In light of Example 5.2, this means that 'most' symbolic dynamical systems admit dimensionally compatible Lipschitz metrics. For example, we have the following result:

Corollary 5.5 *Let* $(\mathbb{V}, \bullet\!\!\to)$ *be a dimensionally homogeneous digraph with a finite estuary* $\mathbb{U}$ (e.g. a biconnected digraph)*. There exists a metric d on $\mathcal{A}^{\mathbb{V}}$ such that, if $\mathcal{X} \subseteq \mathcal{A}^{\mathbb{V}}$ is any pattern space with $\underline{h}_{\mathsf{u}}(\mathcal{X}) > 0$ for some $\mathsf{u} \in \mathbb{U}$, then* $\dim(\mathcal{X}, d) = \dim(\mathbb{V}, \bullet\!\!\to)$*. Furthermore, if $\Phi : \mathcal{A}^{\mathbb{V}} \longrightarrow \mathcal{A}^{\mathbb{V}}$ is a continuous map with network $(\bullet\!\!\to)$, then Φ is d-Lipschitz.*

Proof. Let d be a metric based on $\mathbb{U}$, as in Example 5.2. Then $\overline{\dim}(\mathcal{X}, d) \underset{(*)}{\le} \overline{\dim}(\mathbb{V}, \bullet\!\!\to) \underset{(\dagger)}{=} \dim(\mathbb{V}, \bullet\!\!\to)$, where $(*)$ is by Proposition 5.4(c) and $(\dagger)$ is by dimensional homogeneity. On the other hand, $\underline{\dim}(\mathcal{X}, d) \underset{(*)}{\ge} \underline{\dim}(\mathbb{V}, \bullet\!\!\to) \underset{(\dagger)}{=} \dim(\mathbb{V}, \bullet\!\!\to)$, where $(*)$ is by Proposition 5.4(a) and $(\dagger)$ is by dimensional homogeneity. We conclude that $\dim(\mathbb{V}, \bullet\!\!\to) \le \underline{\dim}(\mathcal{X}, d) \le \overline{\dim}(\mathcal{X}, d) \le \dim(\mathbb{V}, \bullet\!\!\to)$; hence $\dim(\mathcal{X}, d)$ is well-defined and $\dim(\mathcal{X}, d) = \dim(\mathbb{V}, \bullet\!\!\to)$. The fact that d is Φ-lipschitz was demonstrated in Example 5.2. ☐

Example 5.6. (a) Let $\mathbb{V} = \mathbb{Z}^2$ have the Cayley digraph structure induced by generating set $\mathbb{B} := \{(\pm 1, 0)\}, \{(0, \pm 1)\}$. Then $(\mathbb{Z}^2, \bullet\!\!\to)$ is biconnected, so any singleton set is an estuary. So, let $\mathsf{o} = (0, 0)$ be the origin, and let $\mathbb{U} := \{\mathsf{o}\}$; set $c_{\mathsf{o}} = 1$ and $c_{\mathsf{z}} = 0$ for all nonzero $\mathsf{z} \in \mathbb{Z}^2$.

For any $r > 0$, we have $\mathbb{B}(\mathsf{o}, r) := \{\mathsf{z} \in \mathbb{Z}^2 \,;\, |z_1| + |z_2| \le r\}$. Let $\lambda = 2$; then the metric $d_{\mathbf{c},\lambda}$ from Example 5.2 becomes the standard Cantor metric on $\mathcal{A}^{\mathbb{Z}^2}$:

$$d(\mathbf{a}, \mathbf{a}') \quad := \quad \frac{1}{2^R}, \quad \text{where} \quad R \;:=\; \max\left\{r \in \mathbb{N} \,;\, \mathbf{a}_{\mathbb{B}(\mathsf{o},r)} = \mathbf{a}'_{\mathbb{B}(\mathsf{o},r)}\right\}.$$

If $\mathcal{X} \subseteq \mathcal{A}^{\mathbb{Z}^2}$ is any subshift with positive topological entropy, then Proposition 5.4 says $\dim(\mathcal{X}, d) = \dim(\mathbb{Z}^2, \bullet\!\!\to) = 2$. If $\Phi : \mathcal{A}^{\mathbb{Z}^2} \longrightarrow \mathcal{A}^{\mathbb{Z}^2}$ is any cellular automaton whose local rule has neighbourhood $\{0\} \sqcup \mathbb{B}$, then Φ is d-Lipschitz, by Example 5.2.

(b) By a similar argument, if $\mathbb{G}$ is any finitely generated group with growth dimension D and a biconnected Cayley digraph structure, then we can construct a Cantor metric d on $\mathcal{A}^{\mathbb{G}}$ such that, if $\mathcal{X} \subseteq \mathcal{A}^{\mathbb{G}}$ is any subshift with positive topological entropy, then $\dim(\mathcal{X}, d) = D$. Furthermore, if Φ is any CA on $\mathcal{X}$, we can design d to be Φ-Lipschitz.

(c) However, it is possible to construct zero-entropy subshifts of $\mathcal{A}^{\mathbb{Z}^D}$ with dimensions less than D. For example, treat $\mathcal{A}^{\mathbb{Z}^2} \cong \prod_{z\in\mathbb{Z}} \mathcal{A}^{\mathbb{Z}}$ in the obvious way, so that any $\mathbf{a} \in \mathcal{A}^{\mathbb{Z}^2}$ has the form $\mathbf{a} = (\ldots, \mathbf{a}_{-1}, \mathbf{a}_0, \mathbf{a}_1, \mathbf{a}_2, \ldots)$, where $\mathbf{a}_z \in \mathcal{A}^{\mathbb{Z}}$ for all $z \in \mathbb{Z}$. Let $\mathcal{X} := \{(\ldots, \mathbf{a}, \mathbf{a}, \mathbf{a}, \ldots) \; ; \; \mathbf{a} \in \mathcal{A}^{\mathbb{Z}}\} \subset \mathcal{A}^{\mathbb{Z}^2}$. Then $h_{\text{top}}(\mathcal{X}) = 0$ and $\dim(\mathcal{X}, d) = 1$. ◇

Let $(\mathcal{X}, d)$ and $(\mathcal{X}', d')$ be two metric spaces. A continuous function $\Gamma : \mathcal{X} \longrightarrow \mathcal{X}'$ is (d, d')-***Hölder*** if, there exist $\eta, \lambda \in (0, \infty)$ such that, for any $\mathbf{x}_1, \mathbf{x}_2 \in \mathcal{X}$,

$$d'\left(\Phi(\mathbf{x}_1),\ \Phi(\mathbf{x}_2)\right) \quad \leq \quad \lambda \cdot d(\mathbf{x}_1, \mathbf{x}_2)^\eta. \tag{27}$$

For example, any Lipschitz function is Hölder, with $\eta = 1$. If Γ is a homeomorphism, then we say Γ is (d, d')-***biHölder*** if both Γ and Γ^{-1} are Hölder (possibly with different values of η and/or λ).

Proposition 5.7 *Let* $(\mathcal{X}, d)$ *and* $(\mathcal{X}', d')$ *be metric spaces.*

- **(a)** *Let* $\Gamma : \mathcal{X} \longrightarrow \mathcal{X}'$ *be a* (d, d')*-Hölder surjection. Then* $\dim(\mathcal{X}, d) \geq \dim(\mathcal{X}', d')$.
- **(b)** *If* Γ *is a* (d, d')*-biHölder homeomorphism, then* $\dim(\mathcal{X}, d) = \dim(\mathcal{X}', d')$.

Proof. (b) follows from (a). To see (a), suppose Γ is (d, d')-Hölder, and let $\eta, \lambda > 0$ be as in eqn.(27).

Claim 1: *For any* $\epsilon > 0$, $N_\epsilon(\mathcal{X}) \;\geq\; N_{\lambda\epsilon^\eta}(\mathcal{X}')$.

Proof. Let $\mathcal{O} := \{\mathbf{O}_1, \mathbf{O}_2, \ldots, \mathbf{O}_N\}$ be any open ϵ-cover of $\mathcal{X}$. Then for each $n \in [1...N]$, the set $\Phi(\mathbf{O}_n)$ is open (because Φ is an open map, being a continuous surjection onto a compact space), and has diameter at most $\lambda\epsilon^\eta$ by eqn.(27). The collection $\Phi(\mathcal{O}) := \{\Phi(\mathbf{O}_1), \ldots, \Phi(\mathbf{O}_n)\}$ together covers $\mathcal{X}'$, because $\mathcal{O}$ covers $\mathcal{X}$ and Φ is surjective. Thus, $\Phi(\mathcal{O})$ is a $(\lambda\epsilon^\eta)$-diameter open cover of $\mathcal{X}'$. If $\mathcal{O}$ is a minimal open ϵ-cover of $\mathcal{X}$, then $N_\epsilon(\mathcal{X}) = N$. Since $\Phi(\mathcal{O})$ is a $(\lambda\epsilon^\eta)$-cover of $\mathcal{X}'$ with N pieces, we have $N_{\lambda\epsilon^\eta}(\mathcal{X}') \leq N$. ◇ `Claim 1`

It follows that

$$\begin{aligned}
\dim(\mathcal{X}, d) &= \lim_{\epsilon\to 0} \frac{\log[\log(N_\epsilon(\mathcal{X}))]}{\log[-\log(\epsilon)]} \underset{(\ddagger)}{\geq} \lim_{\epsilon\to 0} \frac{\log[\log(N_{\lambda\epsilon^\eta}(\mathcal{X}'))]}{\log[-\log(\epsilon)]} \\
&= \lim_{\epsilon\to 0} \left(\frac{\log[-\log(\lambda\epsilon^\eta)]}{\log[-\log(\epsilon)]}\right) \cdot \left(\frac{\log[\log(N_{\lambda\epsilon^\eta}(\mathcal{X}'))]}{\log[-\log(\lambda\epsilon^\eta)]}\right) \\
&\underset{(*)}{=} \lim_{\epsilon\to 0} \frac{\log[\log(N_{\lambda\epsilon^\eta}(\mathcal{X}'))]}{\log[-\log(\lambda\epsilon^\eta)]} \underset{(\dagger)}{=} \lim_{\epsilon'\to 0} \frac{\log[\log(N_{\epsilon'}(\mathcal{X}'))]}{\log[-\log(\epsilon')]} \quad = \quad \dim(\mathcal{X}', d).
\end{aligned}$$

Here, $(\ddagger)$ is by Claim 1, and $(\dagger)$ is where we make the change of variables $\epsilon' = \lambda\epsilon^\eta$. Finally, $(*)$ is because

$$\begin{aligned}
\lim_{\epsilon\to 0} \frac{\log[-\log(\lambda\epsilon^\eta)]}{\log[-\log(\epsilon)]} &= \lim_{\epsilon\to 0} \frac{\log[-\log(\lambda) - \eta\log(\epsilon)]}{\log[-\log(\epsilon)]} \underset{(\mathrm{H})}{=} \lim_{\epsilon\to 0} \frac{\frac{-\eta/\epsilon}{-\log(\lambda)-\eta\log(\epsilon)}}{\frac{-1/\epsilon}{-\log(\epsilon)}} \\
&= \lim_{\epsilon\to 0} \frac{\eta\log(\epsilon)}{\log(\lambda) + \eta\log(\epsilon)} \quad = \quad 1,
\end{aligned}$$

where (H) is by L'Hospital's rule. □

It follows that the connectivity network dimension of a symbolic dynamical system is invariant under biHölder topological conjugacy.

Corollary 5.8 *Let* $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}_1, \Phi_1)$ *and* $(\mathcal{B}^{\mathbb{W}}, \mathcal{X}_2, \Phi_2)$ *be two symbolic dynamical systems, and let* d_1 *and* d_2 *be dimensionally compatible Lipschitz metrics on* $\mathcal{X}_1$ *and* $\mathcal{X}_2$ *respectively* (e.g. as given by Corollary 5.5).

(a) *If there is a factor mapping* $(\mathcal{X}_1, \Phi_1) \longrightarrow (\mathcal{X}_2, \Phi_2)$ *which is* (d_1, d_2)*-Hölder, then* $\dim(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}_1) \geq \dim(\mathbb{W}, \mathrel{\bullet\!\!\rightarrow}_2)$.

(b) *If* $(\mathcal{X}_1, \Phi_1)$ *and* $(\mathcal{X}_2, \Phi_2)$ *are conjugate via a bi-Hölder homeomorphism, then* $\dim(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}_1) = \dim(\mathbb{W}, \mathrel{\bullet\!\!\rightarrow}_2)$.

Remark 5.9. Clearly, a continuous function $\Phi : \mathcal{A}^{\mathbb{V}} \longrightarrow \mathcal{A}^{\mathbb{V}}$ also admits other Lipschitz metrics which are *not* dimensionally compatible. For example, let $\mathbb{U} = \mathbb{V}$ in Example 5.2. If the coefficient system $\mathbf{c}$ decays slowly enough, we can make $\dim(\mathcal{A}^{\mathbb{V}}, d_{\mathbf{c},\lambda})$ arbitrarily large. However, if $\underline{h}(\mathcal{X}) > 0$, then Proposition 5.4(a) says it is not possible to make $\dim(\mathcal{A}^{\mathbb{V}}, d_{\mathbf{c},\lambda})$ smaller than $\underline{\dim}(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$ for any choice of $\mathbf{c}$. ◊

Conclusion

For any symbolic dynamical system $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$, one can define a digraph structure $(\mathrel{\bullet\!\!\rightarrow})$ on $\mathbb{V}$. We have shown that certain topological-dynamical properties of $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ are related to the connectivity $(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$, and in particular, to its dimension. What other dynamical properties of $(\mathcal{A}^{\mathbb{V}}, \mathcal{X}, \Phi)$ are influenced by the geometry of $(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$?

One could also go the other way. Starting with an infinite digraph $(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$, consider a randomly generated self-map $\Phi : \mathcal{A}^{\mathbb{V}} \longrightarrow \mathcal{A}^{\mathbb{V}}$, such that $(\mathrel{\bullet\!\!\rightarrow})$ is the network of Φ. What are the 'generic' (i.e. almost-certain) properties of $(\mathcal{A}^{\mathbb{V}}, \Phi)$, and how do they depend on the geometry of $(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$?

For example, §3 suggests the following conjecture: *If* $\dim(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}) \leq 1$*, then almost surely,* $(\mathcal{A}^{\mathbb{V}}, \Phi)$ *is equicontinuous. If* $\dim(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow}) > 1$*, then almost surely,* $(\mathcal{A}^{\mathbb{V}}, \Phi)$ *is sensitive.* (The intuition here comes from percolation theory). However, Figure 4 shows that something more than dimension is required; this network has dimension 2, but it has an infinite number of cut points, so a random mapping Φ with this network is almost-surely equicontinuous. Thus, the conjecture above must be augmented with some kind of 'regularity' condition on $(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$.

A closely related question: Suppose we take a system $(\mathcal{A}^{\mathbb{V}}, \Phi)$ and 'mutate' it, by changing the local rule at a small number of vertices. What topological-dynamical properties are 'robust' under such mutations, and how does this depend on the geometry of $(\mathbb{V}, \mathrel{\bullet\!\!\rightarrow})$?

Acknowledgments. This research began during a research leave at Wesleyan University, and was partially supported by the Van Vleck Fund; I am grateful to Ethan Coven for his generous hospitality. This work benefitted from conversations with Ethan, as well as Pierre Tisseur and Reem Yassawi. This research was also supported by NSERC Grant #262620-2008.

References

[BS89] Bohuslav Balcar and Petr Simon. Appendix on general topology. In *Handbook of Boolean algebras, Vol. 3*, pages 1239–1267. North-Holland, Amsterdam, 1989.

[FMM98] Michele Finelli, Giovanni Manzini, and Luciano Margara. Lyapunov exponents versus expansivity and sensitivity in cellular automata. *J. Complexity*, 14(2):210–233, 1998.

[GH00] Yuval Ginosar and Ron Holzman. The majority action on infinite graphs: strings and puppets. *Discrete Math.*, 215(1-3):59–71, 2000.

[Gro99] M. Gromov. Endomorphisms of symbolic algebraic varieties. *J. Eur. Math. Soc. (JEMS)*, 1(2):109–197, 1999.

[IS91] W. Imrich and N. Seifter. A survey on graphs with polynomial growth. *Discrete Math.*, 95(1-3):101–117, 1991. Directions in infinite graph theory and combinatorics (Cambridge, 1989).

[Kůr03] Petr Kůrka. *Topological and symbolic dynamics*, volume 11 of *Cours Spécialisés [Specialized Courses]*. Société Mathématique de France, Paris, 2003.

[McC91] J. McCanna. Fractal dimensions of graphs. In *Proceedings of the Twenty-second Southeastern Conference on Combinatorics, Graph Theory, and Computing (Baton Rouge, LA, 1991)*, volume 84, pages 193–210, 1991.

[Mor95] Gadi Moran. On the period-two-property of the majority operator in infinite graphs. *Trans. Amer. Math. Soc.*, 347(5):1649–1667, 1995.

[MW89] Bojan Mohar and Wolfgang Woess. A survey on spectra of infinite graphs. *Bull. London Math. Soc.*, 21(3):209–234, 1989.

[NR98] Thomas Nowotny and Manfred Requardt. Dimension theory of graphs and networks. *J. Phys. A*, 31(10):2447–2463, 1998.

[NR99] Thomas Nowotny and Manfred Requardt. Pregeometric concepts on graphs and cellular networks as possible models of space-time at the Planck-scale. *Chaos Solitons Fractals*, 10(2-3):469–481, 1999. Superstrings, M, F, S... theory.

[Pet89] Karl Petersen. *Ergodic Theory.* Cambridge University Press, New York, 1989.

[She93] Mark A. Shereshevsky. Expansiveness, entropy and polynomial growth for groups acting on subshifts by automorphisms. *Indag. Math. (N.S.)*, 4(2):203–210, 1993.

[She96] Mark A. Shereshevsky. On continuous actions commuting with actions of positive entropy. *Colloq. Math.*, 70(2):265–269, 1996.

[Tel89] András Telcs. Random walks on graphs, electric networks and fractals. *Probab. Theory Related Fields*, 82(3):435–449, 1989.

[Tel90] András Telcs. Spectra of graphs and fractal dimensions. I. *Probab. Theory Related Fields*, 85(4):489–497, 1990.

[Tel95] András Telcs. Spectra of graphs and fractal dimensions. II. *J. Theoret. Probab.*, 8(1):77–96, 1995.

[Tel01] András Telcs. Fractal dimensions and Martin boundary of graphs. *Studia Sci. Math. Hungar.*, 37(1-2):145–167, 2001.